



## Influence of the extreme weather events on carbon fluxes in southern European taiga

J. Kurbatova, N. Shalukhina, F. Tatarinov, and A. Varlagin

The question on an arrangement, scales and the factors determining ground sink of CO<sub>2</sub> in a forest zone of Russia is opened and discussed. To estimate of global and regional fluxes of carbon for the different periods of time various methods and modeling calculations are used. For an experimental estimation of carbon fluxes for ecosystem level of averaging the eddy covariance method now is widely used. It allows continuously, all-the-year-round, with the high time sanction to register net ecosystem exchange (NEE), fluxes of a water and heat between forest and an atmosphere. Registration of meteorological parameters of an atmosphere is simultaneously carried out. It allows to find the dependence of deposition or issue of carbon on environmental factors. In territory of Russia the observation of carbon fluxes were begun within the framework of the international projects of EU in 1998. Now there is the eddy covariance complex in a southern European taiga, in territory Central Forest Reserve. The measurements are carried out in uneven-age spruce forest (*Sphagnum-Vaccinium myrtillus*). The choice of object of supervision was caused by a wide circulation of similar types of forest in the European taiga.

In general the 10-years period of measurements has captured a wide range of changes of climatic conditions. The years with extreme - droughty and damp vegetative seasons and also years approached to average climatic norms for this region were during the observations. The high daily, seasonal and annual variability of deposition and issue of carbon is characteristic for spruce forests. The results of measurements of NEE have shown that southern taiga can function during the vegetative period both as a source, and as a sink of carbon for an atmosphere. The cumulative fluxes of NEE for the period April - October depend first of all on temperature and precipitation in the spring period with temperature in a range 5-10C and from duration of this period, and for the period of active vegetation, with temperature of air it is more 10C - from humidifying. However for the answer to a question, whether it is formed NEE the next years after years with extreme weather conditions under influence of the current weather conditions, or gas exchange between forest and an atmosphere is consequences of climatic anomalies, demands longer all-the-year-round observation.