

Nuclear planetology: understanding habitable planets as Galactic bulge stellar remnants (black dwarfs) in a Hertzsprung-Russell (HR) diagram

Goetz Roller

Institute for Nuclear Planetology, Geo- and Cosmochronometry, Munich, Germany (goetz.roller@gmail.com)

The Hertzsprung-Russell (HR) diagram is one of the most important diagrams in astronomy. In a HR diagram, the luminosity of stars and/or stellar remnants (white dwarf stars, WD's), relative to the luminosity of the sun, is plotted versus their surface temperatures (T_{eff}). The Earth shows a striking similarity in size (radius ≈ 6.370 km) and T_{eff} of its outer core surface ($T_{eff} \approx 3800$ K at the core-mantle-boundary) with old WD's (radius ≈ 6.300 km) like WD0346+246 ($T_{eff} \approx 3820$ K after ≈ 12.7 Ga [1]), which plot in the HR diagram close to the low-mass extension of the stellar population or main sequence. In the light of nuclear planetology [2], Earth-like planets are regarded as old, down-cooled and differentiated black dwarfs (Fe-C BLD's) after massive decompression, the most important nuclear reactions involved being ⁵⁶Fe(γ , α)⁵²Cr (etc.), possibly responsible for extreme terrestrial glaciations events ("snowball" Earth), together with (γ, n) , (γ, p) and fusion reactions like ¹²C((α, γ) ¹⁶O. The latter reaction might have caused oxidation of the planet from inside out. Nuclear planetology is a new research field, tightly constrained by a coupled ¹⁸⁷Re-²³²Th-²³⁸U systematics. By means of nuclear/quantum physics and taking the theory of relativity into account, it aims at understanding the thermal and chemical evolution of Fe-C BLD's after gravitational contraction (e.g. Mercury) or Fermi-pressure controlled collapse (e.g. Earth) events after massive decompression, leading possibly to an r-process event, towards the end of their cooling period [2]. So far and based upon ¹⁸⁷Re-²³²Th-²³⁸U nuclear geochronometry, the Fe-C BLD hypothesis can successfully explain the global terrestrial MORB $^{232}Th/^{238}$ U signature [3]. Thus, it may help to elucidate the DM (depleted mantle), EMI (enriched mantle 1), EMII (enriched mantle 2) or HIMU (high U/Pb) reservoirs, and the $^{187}Os/^{188}Os$ isotopic dichotomy in Archean magmatic rocks and sediments [4]. Here I present a conceptual model constraining the evolution of a rocky planet like Earth or Mercury from a stellar precursor of the oldest population to a Fe-C BLD, shifting through different spectral classes in a HR diagram after massive decompression and tremendous energy losses. In the light of WD/BLD cosmochronology [1], solar system bodies like Earth, Mercury and Moon are regarded as captured interlopers from the Galactic bulge, Earth and Moon possibly representing remnants of an old binary system. Such a preliminary scenario is supported by similar ages obtained from WD's for the Galactic halo [1] and, independently, by means of 187 Re- 232 Th- 238 U nuclear geochronometry [2, 4, 5], together with recent observations extremely metal-poor stars from the cosmic dawn in the bulge of the Milky Way [6]. This might be further elucidated in the near future by Th/U cosmochronometry based upon a nuclear production ratio Th/U = 0.96 [5] and additionally by means of a newly developed nucleogeochronometric age dating method for stellar spectroscopy, which will be presented in a forthcoming paper. The model shall stimulate geochemical data interpretation from a different perspective to constrain the (thermal) evolution of a habitable planet as to its geo-, bio-, hydro- and atmosphere.

[1] Fontaine et al. (2001), *Public. Astron. Soc. of the Pacific* 113, 409-435. [2] Roller (2015), Abstract T34B-0407, AGU Spring Meeting 2015. [3] Arevalo et al. (2010), *Chem. Geol*. 271, 70-85. [4] Roller (2015), *Geophys. Res. Abstr.* 17, EGU2015-2399. [5] Roller (2015), 78th Annu. Meeting Met. Soc., Abstract #5041. [6] Howes et al. (2015), *Nature* 527, 484-487.