

Modeling and Forecasting the Onset and Duration of a Severe Dutch Fog Event

I.R. van der Velde (1), G.J. Steeneveld (1), B.G.J. Wijchers Schreurs (2), and A.A.M. Holtslag (1)

(1) Wageningen Univ., Meteorology and Air Quality Section, PO box 47, 6700 AA, Wageningen, Netherlands
(ivar.vandervelde@wur.nl), (2) Royal Netherlands Meteorological Institute, PO box 201, 3730 AE, De Bilt, Netherlands

A case of severe radiation fog in the Netherlands is analyzed as a benchmark for the development of a very high resolution NWP model for airport capacity prognoses. Simulations of the mesoscale models WRF and Hirlam are evaluated to determine the state of the art in fog forecasting and to derive requirements for further research and development.

For this case, WRF is unable to simulate the fog for most of the permutations of parameterizations selectable in its framework. Hirlam does model the onset of fog but is unable to let the fog grow beyond the lowest model layer, leading to an early dispersal of fog at the morning transition. The sensitivity of fog forecasts to model formulation is further analyzed with a high resolution single column version of Hirlam, and with an additional single column research model, which was specifically designed for fog forecasting. The single column results are found to be sensitive to the proper specification of initial conditions and external forcings. High vertical resolution is essential for the formation and growth of the fog layer and when the fog lifts for the maintenance of a stratus deck. The properly configured column models are able to accurately model the onset of fog and its maturation, but fail in the simulation of fog persistence and subsequent dispersal. Details of the turbulence parameterization appear to be important in this process. It is concluded that, despite advances in numerical weather prediction, fog forecasting is still a challenge.