

## New insights into the origin of magnetite crystals in ALH84001 carbonate disks

**K.L. Thomas-Keprrta(1), S.J. Clemett(1), D.S. McKay(2) E.K. Gibson(2), S.J. Wentworth(1)**

(1)ESCG at NASA/Johnson Space Center, Houston, TX 77058, (2)KR, ARES, NASA/Johnson Space Center, Houston, TX 77058; (kathie.thomas-keprta-1@nasa.gov)

### Background and Introduction

Martian meteorite ALH84001 preserves evidence of interaction with aqueous fluids while on Mars in the form of microscopic carbonate disks which are believed to have precipitated ~3.9 Ga ago at beginning of the Noachian epoch. Intimately associated within and throughout these carbonate disks are nanocrystal magnetites ( $Fe_3O_4$ ) with unusual chemical and physical properties, whose origins have become the source of considerable debate. One group of hypotheses argues that these  $Fe_3O_4$  are the product of partial thermal decomposition of the host carbonate. Alternatively, the origins of  $Fe_3O_4$  and carbonate may be unrelated; that is, from the perspective of the carbonate the magnetite is allochthonous. We have sought to resolve between these hypotheses through the detailed characterization of the compositional and structural relationships of the carbonate disks and associated magnetites with the orthopyroxene matrix in which they are embedded [1]. We focus this discussion on the composition of ALH84001 magnetites and then compare these observations with those from experimental thermal decomposition studies of sideritic carbonates under a range of plausible geological heating scenarios.

### Methods

*ALH84001.* Seven focused ion beam (FIB) sections were extracted from two carbonate disks -- three from the disk inner cores and four from the thin rims which surround the cores. FIB sections were either micro-welded onto Cu grids or placed onto a continuous C film supported by a Cu grid. Sections were analyzed by high resolution transmission electron microscopy (TEM) equipped with light element energy dispersive X-ray spectroscopy (EDX).

*Roxbury Siderite.* To investigate the chemical compositions of magnetite formed from the decomposition of Fe-rich carbonates, a sample of Roxbury siderite, having a bulk composition of  $(Fe_{0.84}Mg_{0.10}Mn_{0.04}Ca_{0.02})CO_3$  [2], was decomposed using two different heating regimes -- 'slow', ( $\sim 10^{-2}$  K/sec; identical to that used by [3]), and 'fast' ( $\sim 10^8$ - $10^9$  K/sec) using a pulsed  $CO_2$  laser to simulate the rapid heterogeneous heating characteristic of impact

shock. An unheated sample served as a control. Unheated, 'fast' and 'slow' heated samples were embedded in epoxy and prepared for analysis by TEM using diamond knife ultramicrotomy.

### Results

*ALH84001.* Carbonate disks can be envisioned as being composed of three concentric annular zones; starting from the center there is an inner central and outer core surrounded by a thin rim composed of optically alternating black-white-black layers. ALH84001 magnetites are embedded within all compositions of carbonate ranging from the most Fe-rich  $(Fe_{0.60}Mg_{0.16}Ca_{0.05}Mn_{0.04})CO_3$  at the disk core to the most Fe-free magnesite (white layer in the rim zone; see Fig. 1) with the highest population density occurring within the black layers of the rim zone. The majority of these magnetites are stoichiometrically pure  $Fe_3O_4$  although there were several notable exceptions where a small fraction of magnetites contained minor to trace amounts of Cr, which was undetectable in the surrounding host carbonate (Fig.2).

*Roxbury Siderite Control Sample.* Roxbury siderite is compositionally analogous to the Fe-rich component of ALH84001 carbonate disks, with a composition of  $(Fe_{0.750}Mg_{0.240}Ca_{0.004}Mn_{0.003})CO_3$ , as calculated by [4]. However, TEM characterization at the sub-micron scale of Roxbury siderite shows significant variations in Fe:Mg ratio relative to the bulk composition and appeared to be correlated with grain size. The fine grain ( $<100$  nm) component is Mg-poor (Fe:Mg  $> 20:1$ ) while, by comparison, the coarse grained component is Mg-rich (Fe:Mg  $< 8:1$ ). In both size ranges Mn and Ca appeared relatively uniform in comparison.

*Roxbury Siderite 'Fast' and 'Slow' Heated Samples.* Under both 'fast' and 'slow' heating regimes, decomposition of Roxbury siderite resulted in the formation of impure ferrites with the Mg content of the impure ferrite varying from one crystal to another while the Mn content remained invariant. Since the Mg and Mn variation mirrors that of the unheated carbonate in magnitude and range, this variation is

simply a reflection of the initial content of the carbonate (notably, in neither the 'fast' or 'slow' products did we find any evidence for discrete MgO or CaO phases).

These results, in conjunction with prior carbonate decomposition studies relating to the thermochemistry of the Fe-, Mg- and Ca-ternary carbonates (e.g., 5 - 7), show that the decomposition of impure, cation substituted siderites invariably yields impure *Fe*-oxides.

## Discussion & Implications

Based on previous decomposition studies of mixed cation siderites and our decomposition studies of Roxbury siderite, the following summarizes the thermal decomposition reaction of mixed cation siderite and the composition of the resulting product phase:




Although thermal decomposition hypotheses for the observation of chemically pure magnetites in ALH84001 carbonate disks appears, on the surface, to provide a simple inorganic explanation of the observations, it is not applicable to the formation of the vast majority of ALH84001 magnetites. For example, it is difficult to suggest a process by which the magnesite, which is essentially Fe-free, decomposed to form magnetite. Furthermore, it would be difficult to explain the presence of chemically impure magnetites with minor to trace amounts of Cr since this element cannot substitute into the trigonal ( $R\bar{3}c$ ) structure of carbonate.


Thermal decomposition of Roxbury siderite under *both* 'fast' and 'slow' heating resulted in the formation of impure (Mg,Ca,Mn)-ferrites. These findings agree with prior decomposition studies performed under a wide variety of conditions and are in stark contrast to the chemically pure magnetites characteristic of ALH84001 carbonate disks.

The presence of chemically pure ALH84001 magnetite is inconsistent with formation by thermal decomposition of the host carbonate. We suggest that the majority of ALH84001 magnetites has an allochthonous origin and was added to the carbonate system from an outside source. This origin does not exclude the possibility that a fraction is consistent with formation by biogenic processes, as proposed in previous studies.

**Fig.1.** *Left:* TEM view of ALH84001 magnetites (arrows) embedded within the magnesite layer. *Right:* EDX spectra for one of the magnetites (red circle), the surrounding magnesite matrix (blue circle), and the difference spectrum. These spectra show that the matrix is essentially Fe-free while the magnetite is Mg-free. The conclusion is that this magnetite crystal was not formed from decomposition of the host magnesite.



**Fig.2.** *Top views:* FIB section extracted from the core-rim interface of an ALH84001 carbonate disk (left) and high magnification of an elongated magnetite crystal within that region (right). *Bottom views:* Hue-Saturation-Intensity plots for Fe (left) and Cr (right) of the region (top right) showing the elongated ALH84001 magnetite which contains minor Cr.



## References

- [1] Thomas-Keprta, K. et al. (2009) *in press*.
- [2] Lane, M.D. and Christensen, P.R. (1997) *JGR*, 102, 25581-25592.
- [3] Golden, D.C. et al. (2006) *LPSC XXXVII*, Abst #1199.
- [4] Treiman, A.H. (2003) *Astrobiology*, 3, 369-392.
- [5] Jiménez López, C. et al. (2008) *AGU*, Abst. #P51A-1405.
- [6] Gotor, F.J. et al. (2000) *Phys. & Chem. of Min.*, 27, 495-503.
- [7] Gallagher, P.K. and St. J. Warne, S. (1981) *Thermochim. Acta*, 43, 253-267.