



## High sea level for the Last Interglacial: Contribution of the Antarctic ice sheet

G. Lohmann (1,2), X. Wang (1,3), J. Sutter (1), P. Gierz (1), O. Esper (1), R. Gersonde (1), M. Werner (1), M. Pfeiffer (1), J. Müller (1), and A. Rovere (2)

(1) Alfred Wegener Institute, Helmholtz Center for Polar & Marine Research, Bremerhaven, Germany

(gerrit.lohmann@awi.de), (2) University of Bremen, Marum, Bremen, Germany, (3) College of Oceanography, Hohai University, Nanjing, China

Understanding the dynamics of warm climate states has gained increasing importance in the face of anthropogenic climate change. During the Last Interglacial (LIG, ~128 to 116 ka), greenhouse gas concentrations and high latitude insolation were higher than pre-industrial levels, causing a high-latitude warming (Turney and Jones, 2010; Pfeiffer and Lohmann, 2016). As a result of this modestly warmer climate, polar ice sheets were smaller and estimates report that the global mean sea level was 6-9 meters higher than today (Dutton et al., 2015). However, proxy reconstructions indicating a high-stand of LIG sea level are subject to uncertainties in timing and magnitude (Rovere et al. 2016). We present a suite of model results to evaluate the thresholds and feedbacks in the system, and will compare the simulations with paleoclimate reconstructions from high southern latitudes.

Our fully-coupled atmosphere-ocean isotopic simulation of the LIG indicate that temporal and spatial gradients in  $\delta^{18}\text{O}$  do not match, adding uncertainty to the paleothermometer for past warm climates. A simulation using a reduced West Antarctic Ice Sheet (WAIS) is consistent with the isotopic signature found in ice core data (Masson-Delmotte et al., 2011). A reduction of the Greenland Ice Sheet height causes an additional remote warming over the WAIS via atmosphere-ocean teleconnections (Pfeiffer and Lohmann, 2016), adding a destabilizing positive feedback to the system.

Ice sheet model simulations indicate that a pronounced subsurface oceanic warming can destabilize the WAIS, resulting in an oceanic gateway between the Ross and Weddell Seas (Sutter et al. 2016). We detect a threshold behavior of the WAIS in the range of 2-3°C warming. A sensitivity study using the new oceanic gateway between the Atlantic and Pacific Oceans as bathymetrical boundary condition, indicates that this region would be covered by sea ice. Mixing due to sea-ice formation prevents a pronounced warming around the WAIS and would stabilize the WAIS. Thus, the sea level question of the LIG (Sutter et al. 2016; DeConto and Pollard, 2016) is open again.

Past sea-level records located far from Antarctica and hence relatively unaffected by isostatic changes, show that it is possible that the end of the LIG was characterized by a sudden meltwater pulse (O'Leary et al., 2013), that made the sea level rise abruptly. It is possible that a sudden, late-MIS 5e sea level rise was triggered by a collapse of either West or East Antarctica, as it appears that parts of the Greenland Ice Sheet had been already completely melted early in the interglacial. A destabilizing feedback of the WAIS might be related to a pronounced subsurface warming caused by meltwater and a pronounced vertical ocean stratification (Weber et al., 2014).

Keywords: Last Interglacial, Antarctica, sea level, sea ice

### References

1. Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P., Rahmstorf, S., Raymo, M.E., 2015. Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science* (80- ). 349. doi:10.1126/science.aaa4019
2. DeConto, R., Pollard, D., 2016. Contribution of Antarctica to past and future sea-level rise. *Nature*, 531, 591-597.
3. Kopp, R. E., Simmons, F.J., Mitrovica, J.X., Maloof, A.C., Oppenheimer, M., 2009. Probabilistic assessment of sea-level during the last interglacial stage. *Nature*, 462, 836-867.
4. Masson-Delmotte, V., Buiiron, D., Ekaykin, A., Frezzotti, M., Gallée, H., Jouzel, J., Krinner, G., Landais, A.,

Motoyama, H., Oerter, H., Pol, K., Pollard, D., Ritz, C., Schlosser, E., Sime, L. C., Sodemann, H., Stenni, B., Uemura, R., and Vimeux, F.: A comparison of the present and last interglacial periods in six Antarctic ice cores, *Clim. Past*, 7, 397–423, doi:10.5194/cp-7-397-2011, 2011.

5. O’Leary, M.J., Hearty, P.J., Thompson, W.G., Raymo, M.E., Mitrovica, J.X., Webster, J.M., O’Leary, M.J., 2013. Ice sheet collapse following a prolonged period of stable sea level during the last interglacial. *Nat. Geosci.* 6, 796–800. doi:10.1038/ngeo1890

6. Pfeiffer, M. and Lohmann, G., 2016: Greenland Ice Sheet influence on Last Interglacial climate: global sensitivity studies performed with an atmosphere–ocean general circulation model, *Clim. Past*, 12, 1313–1338. doi:10.5194/cp-12-1313-2016

7. Rovere, A., Raymo, M.E., Vacchi, M., Lorscheid, T., Stocchi, P., Gómez-Pujol, L., Harris, D.L., Casella, E., O’Leary, M.J., Hearty, P.J., 2016. The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world. *Earth-Science Rev.* 159, 404–427. doi:10.1016/j.earscirev.2016.06.006

8. Sutter, J., P. Gierz, K. Grosfeld, M. Thoma, and G. Lohmann, 2016: Ocean temperature thresholds for Last Interglacial West Antarctic Ice Sheet collapse. *Geophysical Research Letters*, 43 (6), 2675–2682. doi: 10.1002/2016GL067818

9. Turney, C. S. M. and Jones, R. T.: Does the Agulhas current amplify global temperatures during super-interglacials?, *J. Quaternary Sci.*, 25, 839–843, doi:10.1002/Jqs.1423, 2010.

10. Weber, M. E., Clark, P. U., Kuhn, G., Timmermann, A., Sprenk, D., Gladstone, R., Zhang, X., Lohmann, G., Menziel, L., Chikamoto, M. O., Friedrich, T., Ohlwein, C., 2014: Millennial-scale variability in Antarctic ice-sheet discharge during the last deglaciation. *Nature* 510, 134–138, doi:10.1038/nature13397