



## **TWIRL: Part 2: Very Fine-Scale Dual-Doppler and In Situ Analysis of a Strong Tornado**

Josh Wurman (1), Howard Bluestein (2), Karen Kosiba (3), Dylan Reif (2), Paul Robinson (3), and Zachary Wienhoff (2)

(1) Center for Severe Weather Research, Boulder, CO, United States (jwurman@cswr.org), (2) University of Oklahoma, Norman, OK, United States, (3) Center for Severe Weather Research, Boulder, CO, United States

The OU RAXPOL and CSWR/NSF DOW7 mobile radars deployed near and in front of a large and intense tornado in southern Oklahoma on 09 May 2016, establishing an unprecedentedly short 3.5 km dual-Doppler baseline.

The tornado, exhibiting a complex wind field structure, moved through the dual-Doppler lobes of the radars and crossed between them at a range of  $< 2$  km to each radar. Sweep update rates are 2 s for RAXPOL and 7 s for DOW7, resulting in the shortest time-scale dual-Doppler observations ever in a tornado. Gating is 12.5 m in DOW7 and 75 m for RAXPOL, with  $\sim 1$  degree beam widths. The result is 4D temporal-spatial resolution of 2 s (7 s)  $\times$  27 m  $\times$  27 m  $\times$  75 (12.5) m = 109,000 (64,000) m<sup>3</sup>s for RAXPOL (DOW7) at the tornado's closest approach.

Dual-Doppler analysis with volumetric gridding as small as  $\sim 60$  m  $\times$  60 m  $\times$  60 m, at 7 s intervals, capable of revealing the vector-wind structure of sub-tornadic structures, is possible for the first time. Rotational space-time conversion will be implemented to accurately retrieve features spinning about the tornado. RAXPOL and DOW7 are both dual-polarimetric, DOW7 with two independent dual-polarimetric systems, providing unique fine-scale and intercomparable ZDR and rho-HV observations of tornadic debris lofted by the tornado and its sub-vortices.

RAXPOL and DOW data revealed a large tornado with a core flow diameter of  $\sim 500$  m, exhibiting a complex and rapidly evolving structure including intense sub-vortices. Preliminary dual-Doppler vector wind syntheses, dual-polarimetric analysis will be presented, documenting the evolution of the near-ground wind field and debris structure in this tornado.

This is part 2 of a two-part talk. Kosiba will present the companion talk.