



Identifying water stress in a groundwater dependent ecosystem subjected to groundwater extraction.

R. Froend (1), K. Smettem (2), M. Bertuch (1), C. Xu (3), C. Robertson (3), and M. Martin (3)

(2) University of Western Australia, Centre for Ecohydrology, Environmental Systems Engineering, Perth, Australia

(smettem@sese.uwa.edu.au), (1) Edith Cowan University, Centre for Ecosystem Management, Perth, Australia

(rfroend@ecu.au), (3) Infrastructure Planning Branch, Water Corporation of Western Australia, 629 Newcastle St, Leederville, Perth, W.A. 6007. Australia

Unsympathetic groundwater abstraction practices have had adverse impacts on groundwater dependent ecosystems throughout Australia. In Western Australia, the largest single supply of drinking water for the city of Perth is a superficial aquifer known as the Gnangara mound, located over an area of approximately 2200 km² within and to the north of the city on the Swan coastal plain. Much of the mound is overlain by phreatophytic Banksia woodland that is susceptible to drought stress and death if separated from the unconfined aquifer for prolonged periods over the hot, dry Mediterranean summer. Drought stress has been exacerbated by diminished rainfall due to a changing climate regime.

This paper presents results from a project designed to assess vegetation stress along a transect away from a production bore subsequent to a winter pumping trial. The objective is to determine whether a sustainable pumping regime can be developed that allows water extraction while maintaining a healthy ecosystem.