



Upper Pleistocene climate dynamics from malakofaunal data: records in loess sections of Bansko hill (NE Croatia)

A. Morig (1), O. Mandić (2), D. Pavelić (3), and P. Feric (4)

(1) hgi-cgs, geology department, Sachsova 2, HR-10000 Zagreb, Croatia (adrijano.morig@hgi-cgs.hr), (2) Geological-paleontological Department, Natural History Museum Vienna, Burgring 7, A-1010, Wien, austria, (3) Faculty of Mining and Petroleum Engineering, University of Zagreb, Pierottijeva 6, HR-10000 Zagreb, Croatia, (4) hgi-cgs, geology department, Sachsova 2, HR-10000 Zagreb, Croatia (pavle.feric@hgi-cgs.hr)

Paleontological and taxonomy data from loess section in the Bansko hill (NE Croatia) provide records for climate variations in the upper Pleistocene of SE Europe. Gastropod species from 4 loess sections, suggest a slight dry/humid variations in climate, during upper pleistocene. Periods of dry and cold climate are more often than humid one. Most abundant from 12 species is *Helicopsis striata*, gastropod typical for open and dry habitats. The high relative percentage of dry gastropod taxa in the loess sections, indicates a dry and cold climate for this area, with shorter periods of warmer and more humid climate. This study shows a climate induced changes in taxonomy of gastropods, which is probably expression of 21-ky-precessional cycles.