



Effects of a Corotating Interaction Region on the structure and dynamics of the Saturnian magnetosphere

Nicolas André (1), Philippe Louarn (1), Chris Arridge (2), Laurent Lamy (3), Caitriona Jackman (3), Baptiste Cecconi (4), Andrew Coates (2), Michele Dougherty (3), Dave Young (5), and Philippe Zarka (4)

(1) Centre d'Etude Spatiale des Rayonnements, Solar System, Toulouse, France (nicolas.andre@cesr.fr, 0033 5 61 55 67 01),

(2) Mullard Space Science Laboratory, UK, (3) Imperial College London, UK, (4) Observatoire Paris-Meudon, France, (5) Southwest Research Institute, USA

In order to understand the response of the Saturnian magnetosphere to solar wind dynamic pressure enhancements, we investigate magnetic field and plasma variations observed in-situ by the Cassini spacecraft. We take advantage of two particular orbits (Rev. 26 and 27 in July and August 2006) with similar radial distance, latitude and local time coverage of the Saturnian magnetotail to examine the large-scale structure and dynamics of the nightside current sheet. The observations obtained during these two orbits differ remarkably and some of them present some similarities with the phases of loading/unloading observed during terrestrial substorms. During one of this orbit (Rev. 27), several increases of lobe magnetic pressure are observed, followed by a sudden change in the average position of the current sheet and variations in the longitudinal modulations of the magnetic, plasma and radio fluctuations (short-term dephasing). The deformation of the current sheet lasted for several days before it returned to its past position. Correlation with enhancements in Saturn Kilometric Radiation emissions suggests that solar wind disturbances may have triggered the observed reconfiguration of the Saturnian magnetotail by compressing and relaxing the magnetosphere. We test this hypothesis and provide a plausible interpretation of the observed event that illustrates the Solar-planetary magnetosphere coupling at Saturn.