



Deep-seated gravitational slope deformations in the highest parts of the Czech Flysch Carpathians: predisposition, structure and time constraints

Tomáš Pánek, Veronika Komárková, and Petr Tábořík

Department of Physical Geography and Geoecology, Faculty of Science, University of Ostrava, Ostrava, Czech Republic
(nikakom@centrum.cz / Fax: +420597092323)

The Czech part of the Outer Western Carpathians (OWC) is mainly formed by variously bedded flysch that is susceptible to an occurrence of deep-seated gravitational slope deformations (DSGSDs). On contrary to majority of other World's mountain regions where the DSGSDs have recently been studied, the OWC are generally aseismic, they didn't experience a postglacial deglaciation debuitressing and reach only low to moderate local relief. This study deals with four structurally preconditioned mountain ridges whose ridgetop parts are spectacularly disrupted by the mass movements (Čertův mlýn Mt., Lysá hora Mt., Ondřejník Mt., Smrk Mt.). All these cases are characterized by typical morphological patterns such as tension cracks, double ridges, crevice-type caves, graben-like troughs or uphill-facing scarps. Each study area has been put through a detailed analysis including mapping, speleological research, structural analyses, paleoseismological trenching, 2D electrical resistivity tomography and radiocarbon and OSL dating. All studied DSGSDs are predisposed by mutual interactions between bedding planes, joint sets, and both normal and strike-slip faults within strongly lithologically and tectonically anisotropic flysch massifs. An orientation of gravitational morphostructures well correlates with a structural fabric of the studied region. The extensive geophysical profiling and trenching on the selected sites revealed significantly higher occurrence of the crevice-type caves and other air-filled voids within the anisotropic rock massifs than it was previously stated. Distinctive subsurface zones with extremely high resistivities ($>4000 \text{ ohm.m}$ - indicating caves) often continue outside the morphological expressions of the DSGSDs. This finding indicates that the failures initiate inside the ridges at depths mostly 10-40 m. The paleoseismological trenches applied to three sackung-like features revealed complicated inner structures involving faulted and bended strata, cataclastic bands and opened (partly infilled) crevices. These structures are an evidence of gravitationally activated tectonic elements whereas dislocations evolved by both gravitationally-induced movements (e.g. sackung, lateral spreading etc.) and collapse of rock mass above opened crevices. Bedrock structures are overlaid by both coarse-grained colluvial wedges and fine-grained colluvia. The OSL dating of fine colluvial deposits and the radiocarbon dating of organic deposits infilling the gravitational trenches indicate Holocene age of the studied landforms; some of them evolved even during Late Holocene period. This finding is in contradiction with previous ideas that the massive rock slope failures are mostly of Pleistocene age. Our results indicate that (i) the extensive manifestations of the typical DSGSDs can evolve even in low to medium high mountains - especially, if rock massifs are formed by strongly anisotropic rocks, (ii) beside known processes such as the lateral spreading, toppling and sackung, the important evolution mechanism of the disrupted ridges involves the collapse and the subsidence of the ridgetop zones due to the deep-seated creep-related opening of the crevices, (iii) process of the DSGSDs formation continues within studied region up to recent times by the formation of new gravitational structures and activation of older (Pleistocene-Early Holocene) landforms.

This research was supported by a project from the Grant Agency of the Czech Republic, No. P209/10/0309: "The effect of historical climatic and hydrometeorological extremes on slope and fluvial processes in the Western Beskydy Mts. and their forefield"