



## **Kinetics and reactive transport of N<sub>2</sub>O in a nitrate-contaminated shallow aquifer: How to transfer static Batch Experiments to highly-transient Field Conditions?**

Helmut Geistlinger (1), Daniel Weymann (2), and Reinhard Well (3)

(1) Helmholtz Centre for Environmental Research, Soil Physics, Halle, Germany (helmut.geistlinger@ufz.de), (2) Soil Science of Temperate and Boreal Ecosystems, Büsgen-Institute, University of Göttingen, Büsgenweg 2, 37077 Göttingen, Germany, (3) Institute of Agricultural Climate Research, Johann Heinrich von Thünen Institute, Bundesallee 50, 38116 Braunschweig

N<sub>2</sub>O-production and reduction within the exchange zone are highly variable in space and time and depend on complex reaction rates. The ongoing controversial discussion about the relevance of indirect N<sub>2</sub>O-emission from shallow groundwater has a high degree of uncertainty, because data interpretation is mainly based on regression and on over-simplified steady-state and homogeneous flow models. Moreover prediction models often use non-site specific data sets from literature. For realistic reactive transport modeling of N<sub>2</sub>O we derived a comprehensive site-specific data set of kinetic constants for the depth-dependent heterotrophic and autotrophic denitrification process at the Fuhrberger Feld aquifer. Parameter identification was conducted for incubation experiments using both first-order kinetics and Michaelis-Menten kinetics.

The main question is: How to transfer these kinetic rates obtained from static laboratory experiments to field conditions? For the deeper autotrophic zone we found that the incubation experiments are in good qualitative agreement with field observations, since steady-state flow conditions with nearly constant reducing redox conditions can be assumed. In contrast, N<sub>2</sub>O-production of the heterotrophic exchange zone of shallow groundwater is highly transient related to fluctuating groundwater level (GWL). Therefore, laboratory incubations of aquifer material yielded substantially lower N<sub>2</sub>O concentrations than measured in the field. Thus, the laboratory results are hardly transferable to field conditions. For the first time we develop a reactive transport model that accounts for day-scale changes of the observed GWL and for the short-term fluctuations between anaerobic and aerobic conditions in the exchange zone. We show that the two process time scales (time scale for a typical GWL-fluctuation and time for N<sub>2</sub>O-production) are responsible for N<sub>2</sub>O-production and that under certain conditions N<sub>2</sub>O-accumulation can be observed.

### References

- Geistlinger, H., Ruiyan Jia, D. Eisermann, and C.-F. Stange (2008): Spatial and temporal variability of dissolved nitrous oxide in near-surface groundwater and bubble-mediated mass transfer to the unsaturated zone, *J. Plant Nutrition and Soil Science*, in press.
- Geistlinger, H., G. Wriedt, D. Weymann, and R. Well (2009) Kinetics and reactive transport of N<sub>2</sub>O in a nitrate contaminated shallow aquifer: Development of a process-based reaction module for gas transport, *Vadose Zone Journal*, in review.