



Effects of sulfate aerosols on the East Asian climate in a CGCM.

Won-Mo Kim (1,2), Young Ho Kim (1), Sang-Wook Yeh (1), Byung-Kwon Moon (3), Rokjin Park (2), Daekook Youn (2), Jaemin Jeong (2), Jung-Hun Woo (4), Eul Gyu Im (5), and Chang-Keun Song (6)

(1) Korea Ocean Research & Development Institute, ANSAN, Republic Of Korea, (2) School of Earth and Environmental Sciences, Seoul National University, Seoul 151-747, Korea, (3) Division of Science Education/Institute of Fusion Science, Chonbuk National University, Jeonju, Korea, (4) Department of Advanced Technology Fusion, Konkuk University, Seoul, Korea, (5) Department of Computer Science and Engineering, Hanyang University, Seoul, Korea, (6) Climate Change Research Division, National Institute of Environmental Research, Incheon, Korea

An increase of anthropogenic greenhouse gases and atmospheric aerosols due to human activity is the main forcing responsible for a climate change over the globe. While effects of anthropogenic greenhouse gases have been studied extensively, a role of atmospheric aerosols is highly uncertain so far. Therefore, it is important to understand how aerosol forcing is associated with climate variability to assess their impacts on climate change. In this study, we performed a number of simulations using a Community Climate System Model (CCSM) version 2.0.1 to identify a role of sulfate aerosol forcing on East Asian climate. It is found that a regional aerosol forcing (i.e., sulfate aerosol) could modify the rainfall variability over northeast Asia during spring and summer time. Based on a long term period of simulation, in addition, we could find that sulfate aerosol forcing is associated with changes in the North Pacific Ocean SST variability.