



Melt localization and its relation to deformation in the subcontinental mantle: a case study from layered dunite-harzburgite-lherzolite bodies of the Ronda peridotite massif, Spain

Karoly Hidas (1), Carlos J. Garrido (1), Jean-Louis Bodinier (2), Andrea Tommasi (2), Guillermo Booth-Rea (3), Fernando Gervilla (1), and Claudio Marchesi (2)

(1) Instituto Andaluz de Ciencias de la Tierra, CSIC-UGR, Granada, Spain (karoly.hidas@gmail.com), (2) Geosciences Montpellier, CNRS-Université de Montpellier-2, Montpellier, France, (3) Department of Geodynamics, Universidad de Granada, Granada, Spain

The processes that take place during the transport of melts through the convecting mantle are the least understood and, therefore, state-of-the art problems among a series of processes of formation and evolution of mantle magmas. It is widely accepted that, dunite channels might be pathways by which mantle melts easily pass through the overlying mantle (e.g. Kelemen et al., 1997). The role of shear strain during the formation of dunite bodies in ophiolites was considered in details by Kelemen & Dick (1995). It was also shown that the stress field can control the melt migration paths marked by dunite bodies occurring oriented regularly relative to the hinge and axial plane of a harzburgite fold (Savelieva et al., 2008). The localization of melt flow and formation of channels under mechanical instability during the formation of dunites is expected to lead to a stronger olivine crystallographic preferred orientation (CPO) in these rocks than in their surroundings. However, accepted models explain formation of dunitic lithology mostly in oceanic environment, but one would face several challenges trying to apply them to the subcontinental lithospheric mantle.

The Ronda massif (southern Spain) is the largest (ca. 300km²) of several orogenic peridotite massifs exposed in the Betic and Rif (northern Morocco) mountain belts in the westernmost part of the Alpine orogen that was tectonically emplaced during early Miocene times. One of the most remarkable features of the Ronda massif is the 'recrystallization front' that represents the transition from the spinel-tectonite to the coarse granular peridotite domain corresponding to a narrow boundary of a partial melting domain caused by thinning and coeval asthenospheric upwelling formed at the expense of former subcontinental lithospheric mantle and associated with melting and kilometer-scale migration of melts by diffuse porous flow through the 'asthenospherized' domain (Van der Wal & Bodinier, 1996; Lenoir et al., 2001; Vauchez & Garrido, 2001). In the vicinity of the recrystallization front, coarse granular peridotites pass into layered granular peridotites with a typical layered structure composed of plagioclase lherzolites, harzburgites and dunites.

The main scientific goals of this study are to test new mechanism(s) for the formation of dunites and dunite-harzburgite-lherzolite layered bodies in the subcontinental lithospheric mantle on the example of Ronda peridotite massif (Spain), and to introduce new processes that are expected to lead the evolution of the subcontinental lithospheric mantle in extensional settings.

References

- Kelemen, P.B. & Dick, H.J.B. (1995). Focused melt flow and localized deformation in the upper-mantle — juxtaposition of replacive dunite and ductile shear zones in the Josephine peridotite, SW Oregon. *J. Geophys. Res.* 100 (B1): 423–438.
- Kelemen, P.B., Hirth, G., Shimizu, N., Spiegelman, M. & Dick, H.J.B. (1997). A review of melt migration processes in the adiabatically upwelling mantle beneath oceanic spreading ridges. *Philosophical Transactions of the Royal Society a-Mathematical Physical and Engineering Sciences* 355(1723): 283-318.
- Lenoir, X., Garrido, C.J., Bodinier, J.L., Dautria, J.M. & Gervilla, F. (2001). The recrystallization front of the Ronda peridotite: Evidence for melting and thermal erosion of subcontinental lithospheric mantle beneath the Alboran basin. pp. 141-158.

Savelieva, G.N., Sobolev, A.V., Batanova, V.G., Suslov, P.V. & Bruhmann, G. (2008). The structure of melt transport mantle channels. *Geotectonics* 42 (6): 430—447.

Van Der Wal, D. & Bodinier, J.L. (1996). Origin of the recrystallisation front in the Ronda peridotite by km-scale pervasive porous melt flow. *Contributions to Mineralogy and Petrology*, 122(4): 387-405.

Vauchez, A. & Garrido, C.J. (2001). Seismic properties of an asthenospherized lithospheric mantle: constraints from lattice preferred orientations in peridotite from the Ronda massif. *Earth and Planetary Science Letters* 192(2): 235-249.