



Seasonal and inter-annual variations of community metabolism rates of a *Posidonia oceanica* seagrass meadow

Willy Champenois (1), Gilles Lepoint (2), Bruno Delille (1), Marc-Vincent Commarieu (1), Marilaure Grégoire (2), Jean-Marie Beckers (3), and Alberto V. Borges (1)

(1) University of Liège, Institut de Physique (B5), Chemical Oceanography Unit, Liège, Belgium (alberto.borges@ulg.ac.be, +32-(0)4-3663367), (2) Oceanology Laboratory, University of Liège, Belgium, (3) G.H.E.R., University of Liège, Belgium

We report gross primary production (GPP), community respiration (CR) and net community production (NCP) over *Posidonia oceanica* meadow at 10 m in Corsica (Bay of Revellata) based on the open water O₂ mass balance approach from a data-set of hourly measurements with an array of three O₂ optodes that was deployed from August 2006 to October 2009. The method was checked by the comparison with discrete measurements of metabolic rates derived from benthic chamber incubations also based on the diel change of O₂. This comparison was satisfactory and actually highlights the potential caveats of benthic incubation measurements related to O₂ accumulation in small sized chambers leading to photorespiration, and an under-estimation of GPP. Our data-set confirms previously established knowledge on *P. oceanica* meadows community metabolism: these communities were characterized by intense GPP and CR values, with strong seasonal variations, and were net autotrophic at annual scale. However, the high resolution data-set we obtained reveals additional knowledge that was missed by discrete benthic measurements with a coarse resolution (at best monthly). There was a strong day-to-day variability of GPP and CR, linked to changes in light availability, and extremely high but transient GPP events were recorded. Strong inter-annual variability of community metabolic rates was evidenced, with lower GPP and NCP during the 2006/2007 yearly cycle characterized by a mild and less stormy winter compared to the 2007/2008 and 2008/2009 yearly cycles. This finding suggests that one possible future evolution of carbon flows in *P. oceanica* meadows would be the decrease of export organic carbon to adjacent communities and a decrease of GPP and NCP, since a decrease in frequency and intensity of marine storms is expected in future in the Mediterranean Sea, due to a northward shift of the Atlantic storm track.