



## **Biogeochemistry and carbon mass balance of a coccolithophore bloom in the northern Bay of Biscay (June 2006)**

Jérôme Harlay (1,2), Lei Chou (2), Caroline De Bodt (2), Nicolas Van Oostende (3), Judith Piontek (4), Anja Engel (4), Koen Sabbe (3), Steve Groom (5), and Alberto V. Borges (1)

(1) University of Liège, Institut de Physique (B5), Chemical Oceanography Unit, Liège, Belgium (alberto.borges@ulg.ac.be, +32-(0)4-3663367), (2) Laboratoire d'Océanographie Chimique et Géochemie des Eaux, Université Libre de Bruxelles, Campus de la Plaine, CP208, boulevard du Triomphe, B-1050 Brussels, Belgium, (3) Protistology and Aquatic Ecology, Gent University, Krijgslaan 281-S8, B-9000 Gent, Belgium, (4) HGF Young Investigators Group, Global change and the future marine carbon cycle, Alfred Wegener Institute, Am Handelshafen 12, D-27570 Bremerhaven, Germany, (5) Remote Sensing Group, Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, United Kingdom

Primary production (PP), calcification (CAL), bacterial production (BP) and dark community respiration (DCR) were measured along with a set of various biogeochemical variables, in early June 2006, at several stations at the shelf break of the northern Bay of Biscay. The cruise was carried out after the main spring diatom bloom that, based on the analysis of a time-series of remotely sensed chlorophyll-a (Chl-a), peaked in mid-April. Remotely sensed sea surface temperature (SST) indicated the occurrence of enhanced vertical mixing (due to internal tides) at the continental slope, while adjacent waters on the continental shelf were stratified, as confirmed by vertical profiles of temperature acquired during the cruise. The surface layer of the stratified water masses (on the continental shelf) was depleted of inorganic nutrients. Dissolved silicate (DSi) levels probably did not allow significant diatom development. We hypothesize that mixing at the continental slope allowed the injection of inorganic nutrients that triggered the blooming of mixed phytoplanktonic communities dominated by coccolithophores (*Emiliania huxleyi*) that were favoured with regards to diatoms due to the low DSi levels. Based on this conceptual frame, we used an indicator of vertical stratification to classify the different sampled stations, and to reconstruct the possible evolution of the bloom from the onset at the continental slope (triggered by vertical mixing) through its development as the water mass was advected on-shelf and stratified. We also established a carbon mass balance at each station by integrating in the photic layer PP, CAL and DCR. This allowed computation at each station of the contribution of PP, CAL and DCR to CO<sub>2</sub> fluxes in the photic layer, and how they changed from one station to another along the sequence of bloom development (as traced by the stratification indicator). This also showed a shift from net autotrophy to net heterotrophy as the water mass aged (stratified), and suggested the importance of extracellular production of carbon to sustain the bacterial demand in the photic and aphotic layers.