



## **Intermontane Basin and Landscape Evolution in the Eastern Cordillera of NW Argentina – S Humahuaca Basin (23-24° S)**

Heiko Pingel (1), Strecker Manfred R. (1), Hilley George E. (2), Alonso Ricardo N. (3), and Schmitt Axel K. (4)

(1) Institut für Erd- und Umweltwissenschaften, Universität Potsdam, Potsdam, Germany (heikopingel@googlemail.com),

(2) Department of Geological and Environmental Sciences, Stanford University, Stanford, USA, (3) Facultad de Ciencias

Naturales, Universidad Nacional de Salta, Salta, Argentina, (4) Department of Earth and Space Sciences, UCLA, Los Angeles, USA

Intermontane basins are important repositories for sediment storage, documenting the tectonic, climatic, and sedimentary evolution of orogens in time and space. The Quebrada de Humahuaca in the Southern Central Andes of NW Argentina is an ideal setting to study the evolution of an intermontane depositional environment, which is located within a very narrow section of the Eastern Cordillera between 23-24°S. Elevated due to progressive shortening and associated surface uplift during ongoing orogeny, accompanied by successive aridification due to orographic barrier uplift, this basin has accommodated and retained a variety of Neogene to Quaternary deposits, documenting the late Cenozoic basin evolution, e.g., the transition from an open foreland to an intermontane basin environment. Here, we present new U-Pb zircon ages and paleocurrent reconstructions, suggesting that prior to ~4.3 Ma, the Quebrada de Humahuaca was an integral part of a contiguous foreland basin, whose fluvial network was subsequently decoupled from the foreland due to basement-block uplifts in the course of an eastward migrating deformation front. Subsequently, the basin experienced at least two cycles of hydrologic isolation and re-capture, resulting in repeated filling and re-excavation. The ultimate basin-filling episode between ~1 Ma and <40 ka was followed by episodic incision, leading to abandoned fluvial terraces and pediments >350 m above the present-day base level. This seems temporally consistent with the latest basin-fill history from the neighboring Toro basin and may be related to superposed, climate-driven processes that affected intermontane basins in NW Argentina on a regional level. Renewed evidence for faulting, regional unconformities, and deformed landforms suggest a reactivation of basin-bounding faults and structures within the basin. Detailed observations regarding sedimentary lithologies reveal a pattern of out-of-sequence deformation within the basin that may be associated with the different cycles of alluviation and basin evacuation.