



Water table change differently affects CO₂ and N₂O fluxes in a bio-energy poplar plantation

Donatella Zona, Ivan Janssens, Melanie Verlinden, Laura Broeckx, Joris Cools, and Reinhart Ceulemans
University of Antwerp, Belgium (donatella.zona@ua.ac.be)

We are continuously monitoring O₃, CH₄, CO₂, H₂O and N₂O fluxes from a fast-growing high-density poplar (*Populus*) plantation with eddy covariance from June 2010. In this contribution we present a selection of the data from the first field season when an intense precipitation event (~80 mm rainfall in 48 hours) occurred after a prolonged fairly dry summer period. This first extreme precipitation caused peak N₂O emissions (up to 2200 $\mu\text{g N}_2\text{O-N m}^{-2} \text{h}^{-1}$). However successive rainfall events and similar soil moisture and water table fluctuations did not lead to N₂O emissions of the same magnitude of these first peak emissions, probably because of depletion of the soil nitrogen substrate. In contrast, CO₂ fluxes, both net ecosystem exchange (NEE) and ecosystem respiration (ER) did not respond to any of these rain events, contrary to what has been observed for various other ecosystems. This was probably caused by the N availability to microorganisms that exceeded C availability at our site. Overall the data presented provide important insights in the complexity of the environmental controls on CO₂ and N₂O emission, and the variability in their response to hydrological changes. We present some of the preliminary analysis of the water table and soil moisture effects on CO₂ and N₂O fluxes, and we quantify the overall global warming potential, GWP, of the first growing season for this highly productive ecosystem.