



First retrievals of carbon tetrafluoride (CF₄) from ground-based FTIR measurements: production and analysis of the two-decadal time series above the Jungfrauoch

Pierre Duchatelet (1), Rodolphe Zander (1), Emmanuel Mahieu (1), Jens Mühle (2), Philippe Demoulin (1), Bernard Lejeune (1), Ginette Roland (1), Christian Servais (1), and Olivier Flock (1)

(1) University of Liège, Institute of Astrophysics and Geophysics, Liège, Belgium (p.duchatelet@ulg.ac.be), (2) Scripps Institution of Oceanography, University of California, San Diego, USA

Carbon tetrafluoride (CF₄ or PFC-14) is a potent greenhouse gas that is almost 7400 times more effective (100-yr horizon) than CO₂ on a per molecule basis (IPCC, 2007). This high global warming potential, coming from its medium absorbance combined with a very long atmospheric lifetime (>50000 years; Ravishankara et al., 1993), makes CF₄ a key species among the various greenhouse gases targeted by the Kyoto Protocol. In the Northern hemisphere, current atmospheric CF₄ concentrations are close to 78 pptv, with a large fraction (around 35 pptv, Mühle et al., 2010) coming from natural processes like lithospheric emissions (Harnisch and Eisenhauer, 1998). In addition, CF₄ has been used increasingly since the eighties in electronic and semiconductors industry. The primary aluminum production processes have also been clearly identified as an important anthropogenic source of CF₄ emissions. The partitioning between these two main sources is however problematic, principally due to lacking or incomplete CF₄ emission factors from inventories performed in industrial fields (e.g. International Aluminum Institute, 2009).

Recent in situ ground level measurements of CF₄ in the Northern hemisphere (Khalil et al., 2003; Mühle et al., 2010) or remotely from space (Rinsland et al., 2006) have indicated a significant slowdown in the increase rate of atmospheric CF₄. This probably results from measures adopted by the aluminum industry aiming at the reduction of the frequency and duration of “anode effects” and therefore of related PFCs emissions (International Aluminum Institute, 2009).

The present contribution reports on the long-term evolution (1990-2010) of the atmospheric carbon tetrafluoride total vertical abundance derived from ground-based Fourier transform infrared (FTIR) solar spectroscopy observations around 1285 cm⁻¹ at the Jungfrauoch (46.5°N, 8.0°E, 3580m asl) and compares our findings with results available in the literature. To our knowledge, no equivalent time series (i.e. based on ground-based FTIR technique) has been published to date.

Acknowledgements:

The University of Liège contribution to present work has primarily been supported by the SSD and PRODEX programs funded by the Belgian Federal Science Policy Office, Brussels. We thank the International Foundation High Altitude Research Stations Jungfrauoch and Gornergrat (HFSJG, Bern) for supporting the facilities needed to perform the observations.

References:

Harnisch, J. and A. Eisenhauer, Natural CF₄ and SF₆ on Earth, *Geophys. Res. Lett.*, 25, 2401-2404, 1998.

International Aluminum Institute, The International Aluminum Institute Report on the Aluminum Industry's Global Perfluorocarbon Gas Emissions Reduction Programme – Results of the 2007 Anode Effect Survey, International Aluminum Institute, London, 2009.

IPCC, Climate change 2007: The scientific basis, Tech. rep., Intergovernmental Panel on Climate Change, IPCC/WMO/UNEP, 2007.

Khalil, M. A. K., R. A. Rasmussen, J. A. Culbertson, et al., Atmospheric perfluorocarbons, *Environ. Sci. Technol.*, 37, 4358-4361, 2003.

Möhle, J., A. L. Ganesan, B. R. Miller, et al., Perfluorocarbons in the global atmosphere: tetrafluoromethane, hexafluoromethane, and octafluoropropane, *Atmos. Chem. Phys.*, 10, 5145-5164, 2010.

Ravishankara, A. R., S. Solomon, A. A. Turnispeed and R. F. Warren, Atmospheric lifetimes of long-lived halogenated species, *Science*, 259, 194-199, 1993.

Rinsland, C. P., E. Mahieu, R. Zander, et al., Long-term stratospheric carbon tetrafluoride (CF₄) increase inferred from 1985-2004 space-based solar occultation measurements, *Geophys. Res. Lett.*, 33, L02808, doi:10.1029/2005GL024709, 2006.