



Paraglacial processes operating on High Arctic coastal margins - recent advances from Svalbard.

Matt Strzelecki (1,2)

(1) Department of Geography, Durham University, Durham, UK, (2) Faculty of Geosciences, Adam Mickiewicz University, Poznan, Poland

In contrast to mid and low latitude coasts, relatively little is known regarding the potential impacts of climate and sea-level change on high latitude coastal margins. Indeed, many of the existing intellectual paradigms regarding the functioning of polar beaches are now out-dated, based on descriptive geomorphology and a limited process-based understanding.

My research project aims to address this deficiency in understanding by quantifying the mechanisms and patterns of recent adjustment of High Arctic coasts in Svalbard following the end of Little Ice Age, an interval of time characterized by abrupt climate warming, rapid deglaciation and rising relative sea-levels. The work involves a combination of field based survey, conducted in summer and winter over several seasons, as well as remotely sensed data that uses time series analysis of remotely sensed data from the XX century.

The preliminary results document dramatic changes in sediment flux and coastal response under an intervals characterised by a warming climate, retreating local ice masses, a shortened winter sea-ice season and melting permafrost. These (largely) terrestrial processes are interacting with an upwards trend in relative sea-level attributed to glacio-isostatic land subsidence and on-going global sea-level rise. The pristine coasts of Svalbard provide a superb opportunity to quantify how Arctic coasts are responding to rapid climate warming.

In this paper, I summarise my PhD research to date by presenting results from an analysis of digital aerial photogrammetry, combined with field-based geomorphological mapping. The geographical focus is Petunia Bay, one of the most protected bays of the Svalbard Archipelago, which is characterized by a semi-arid, sub-polar climate, limited wave fetch and tidal range, and rapid retreat rate of all surrounding glaciers. Landscape and coastal change is mapped using combination of DEM-s developed over a 70 year period between 1936 and 2009, combined with field-based geomorphological mapping. I also detail the characteristics of beach sediment from several beach settings together with 'Jennings and Shulmeister' beach classification methods to identify potential factors controlling the morphodynamics of these polar beaches.

This work highlights the need for a greater understanding of the controls on polar coastal sediment budgets, especially given the potential for accelerated warming and sea-level rise in the coming decades and centuries.

This work is funded by the Polish Ministry of Science and Higher Education (grant no. N306 284335).

M. Strzelecki is supported by the Crescendum Est Polonia fellowship and AMU Foundation scholarship