


Microbiology in practice: Are there really any live and active bacteria cultures in yogurts? - a methodical contribution -

Simona Šafarčíková, e-mail: safarcicka@email.cz

THE ANGLO-CZECH HIGH SCHOOL, 1010 Trebizskeho, Ceske Budejovice 5, CZ 370 06, CZECH REPUBLIC, e-mail: cag@cag.cz, http://www.cag.cz



Material

and Methods

START

Publicity

publishing

news

on the school

website

Discussion

Introduction

- Yogurts are a common part of our everyday diet. Most of them are claimed to contain probiotic bacteria cultures. Among the most common bacteria present in yogurt cultures belong so-called lactic acid bacteria, e.g. genus Lactobacillus and Bifidobacterium, which are well known for their positive influence on the intestinal microflora and even for their potential antitumor activity (Gomes & Malcata 1999).
- Since recently rumours have been circulating that yogurts actually do not contain any active bacteria cultures, students are wondering if consuming yogurts is meaningful or not. Thus, they carry out a microbiological experiment to test the presence of probiotic bacteria in yogurts.
- As the poster is intended especially as a methodical contribution, most of the poster area is assigned to the section Material and Methods. Only brief selection of students' results together with short discussion and conclusion is shown.

Aims 1) to find out if there are live and active bacteria cultures in yogurts; 2) to study if there is any kind of biological contamination in yogurts

Information survey

 searching for and studying information sources on lactic acid bacteria and yogurts

the yogurt supply **List of yogurts** in supermarkets

 discussing and setting up the list of yogurts to be tested



Inoculation

- inoculating yogurt cultures on the agar plate on the Petri dish with the inoculation loop
- preparing three replications of each yogurt sample

Cultivation

 putting the Petri dish upside down and cultivating it at 20 °C for 5-7

 assessing the presence and viability of lactic acid bacteria on the scale 1-3 (level 3 is the most viable culture)

Evaluation

 assessing the presence and type of contamination (yeast, mould, other bacteria)

Viability of lactic acid bacteria culture in tested jogurts


Yogurt supply

screening

and specialized

health food


shops



■ absent/non-viable culture hardly viable culture medium-viable culture very viable culture

Results

Biological contamination of tested jogurts no contamination



moulds and bacteria moulds yeasts

bacteria

Fig. 1 Viability of lactic acid bacteria culture in 11 different yogurts (N=3).

Presentation

presenting the experiment and its results

Research report

 compiling a research report (introduction, material and methods, results, discussion, conclusion, references)



Data processing

 transforming the obtained data into the visual form (graphs)

Determination

Fig. 2 Presence and types of biological contamination of 11 different yogurts (N=3).

- determining present genera of lactic acid bacteria, possibly of yeasts, moulds and other bacteria
- Academy of Sciences

- cooperation with microbiologists from the Institute of Soil Biology, Biology Centre of the

Conclusion and future plans

Results may be **skewed**, e.g. because of the lack of the strictly sterile environment at the school lab. Thus, the contamination of yogurts may originate from somewhere else than from yogurts themselves. This practical enables students to use microbiology in practise and to obtain surprising results concerning their everyday diet. In the future, avoiding contamination from the outside and higher number of replication for each sample are planned. The later is essential be able to test the results statistically to make them more reliable.

