



High preindustrial $\delta^{13}\text{C}$ in the deep Arctic Ocean: Implications for the interpretation of abrupt benthic $\delta^{13}\text{C}$ excursions during the last glaciation

A. Mackensen

Alfred Wegener Institute, Bremerhaven, Germany (andreas.mackensen@awi.de)

Only recently a strong air-sea signature of $\delta^{13}\text{C}$, i.e. $\delta^{13}\text{C}_{\text{as}}$, in modern bottom waters of the Atlantic sector of the Antarctic Ocean was revealed (Mackensen, 2012). This high southern $\delta^{13}\text{C}_{\text{as}}$ may have been reduced in the Last Glacial Maximum, and parts of an ocean wide deep-water $\delta^{13}\text{C}_{\text{DIC}}$ lowering may be attributed to sea-ice formation with low- $\delta^{13}\text{C}_{\text{as}}$ brine rejection and diminished air-sea gas exchange in the southern ocean. Low benthic $\delta^{13}\text{C}$ values from the Nordic seas in cold stadials of the last glaciation have been attributed to brine formation, but little is known about the carbon isotopic composition of Arctic Ocean brines and deep-water masses. Here I show that today dissolved inorganic carbon (DIC) of bottom water in the deep Arctic Ocean is ^{13}C enriched with mean $\delta^{13}\text{C}_{\text{DIC}}$ values of 1.2 ‰ whereas bottom waters bathing most of the continental margins with mean $\delta^{13}\text{C}_{\text{DIC}}$ values of about 0.8 ‰. This difference is also recorded in Recent epibenthic foraminiferal $\delta^{13}\text{C}$ from the deep Arctic versus Greenland and Svalbard continental margins. It is in contrast, however, to the continental slope of the Laptev and East Siberian seas, where epibenthic $\delta^{13}\text{C}$ is as high as in the deep basins. I conclude that (i) most of the shelves contributing to Arctic bottom water by brine rejection produce high- $\delta^{13}\text{C}_{\text{as}}$ brine, and (ii) a strong $\delta^{13}\text{C}_{\text{as}}$ signal from brine formation in polynyas today is masked by anthropogenically lowered atmospheric $\delta^{13}\text{C}_{\text{CO}_2}$. I then hypothesize that during stadials, when most of the Arctic Ocean was perennially sea-ice covered, less brine was produced, and that this cessation of brine rejection would have lowered bottom-water $\delta^{13}\text{C}$ values in the Arctic Ocean and subsequently in the Nordic seas.