



## Fracturing and carbonate mineralization in Palaeogene carbonate rocks from the United Arab Emirates: A record of fluid flow

Ihsan Al-Aasm (1), Manhal Sirat (2), Sadoon Morad (3), Ala AlDahan (4), and Osama Al-Jallad (5)

(1) University of Windsor, Earth and Environmental Sciences, Windsor, Canada (alaasm@uwindsor.ca), (2) ADCO, P.O. Box 270, Abu Dhabi, UAE, (3) Department of Petroleum Geosciences, the Petroleum Institute, P.O. Box 2533, Abu Dhabi, United Arab Emirates, (4) Department of Earth Sciences, Uppsala University, 752 36 Uppsala, Sweden, (5) Abu Dhabi Drilling Co, Abu Dhabi

Fracture measurements, field observations, petrographic examination, fracture and geochemical analyses, and fluid-inclusion microthermometry of vein calcite and dolomite helped to unravel the origin of the fracture systems and geochemical evolution of diagenetic fluids, which circulated in Eocene marine carbonates of Jabal Hafit anticline, United Arab Emirates. The vein carbonates consist of coarse-crystalline calcite (mm to 7 cm in diameter) with diverse crystal shapes (equant blocky, bladed and fibrous) and saddle dolomite embedded in limestone and dolostone host rocks.

Two fracture systems have been identified with relatively different time of origin; each of which comprises an extensional set and two conjugate shear sets, wherein the flux of the fluids occurred during tectonic activities of compression and shear related to the Zagros orogeny (Eocene-present).

The  $[U+O]$  18OV-PDB values of vein calcite ( $-20.2\text{\textperthousand}$  to  $-2.9\text{\textperthousand}$  coupled with fluid-inclusion microthermometry (homogenization temperatures  $100^\circ\text{C}$  to  $190^\circ\text{C}$  and salinity of 3-29 wt.% equivalent NaCl) suggest precipitation from hydrothermal fluids of variable salinities and probably origins. In contrast,  $[U+O]$  18OV-PDB values of saddle dolomite in fractures and fossil molds ( $-7.16$  to  $-6.22\text{\textperthousand}$  show less depleted values than their calcitic counterparts but their fluid-inclusion microthermometry (homogenization temperatures  $91.3$  to  $97^\circ\text{C}$  and salinity of 20.5 to 22.0 wt. % NaCl) show more saline fluids than the ones precipitated fracture-filling calcite cements. The correlation between the  $[U+O]$  18OV-PDB and  $[U+O]$  13CV-PDB values in calcite and dolomite reflects two trends; a temperature fractionation dependent trend and a mixing fluid trend. In the latter the possibility of input of dissolved carbon derived from the host carbonate rocks and from meteoric waters may have occurred during tectonic uplift. Strontium isotope ratios of many vein calcite (0.70779-0.70875) and saddle dolomite (average 0.70782) are similar to Palaeogene seawater, and thus are suggested to be derived from the interaction of fluids with the host carbonate rocks. However, radiogenic Sr ratios of some vein calcite and dolomite suggest influx of basinal fluids at high temperature and elevated salinity.