



K-Ar Geochronology of Igneous Amphibole Macrocrystals of Miocene to Pliocene Volcaniclastics, Styrian Basin, Austria

Hans-Peter Bojar (1), Ana-Voica Bojar (2), Stanisław Hałas (3), and Arthur Wojtowicz (3)

(1) Department of Earth Sciences - Mineralogy, Studienzentrum Naturkunde, Universalmuseum Joanneum, Weinzötlstraße 16, A-8045 Graz, Austria, (2) Department of Geology, Salzburg University, A-5020 Salzburg and Department of Mineralogy, Universalmuseum, A-8045 Graz, Austria, (3) Mass Spectrometry Laboratory, Institute of Physics, Maria Curie-Skłodowska University, PL-20-031 Lublin, Poland

The Styrian Basin basin is situated at the south-eastern margin of the Eastern Alps. Two basement highs separate the Styrian Basin in sub-basins: the Middle Styrian Basement High divides the shallower Western Styrian Basin (with \sim 800 m thick Neogene sediments) from the Eastern Styrian Basin (with a \sim 3000 m thick infill). A second, the South Burgenland Basement High, separates the West Pannonian from the Styrian Basin. The north-south trending Auersbach Basement High separates the Eastern Styrian basin in two sub-basins.

The Eastern Styrian Basin hosts distinct volcanic depocentres represented by Miocene (Karpatian to Badenian) shield volcanoes and volcanic tuff intercalations situated within Badenian sediments. A second phase of basaltic volcanism has a Late Miocene to Pliocene age. The Western Styrian Basin hosts only one Middle Miocene volcano. Most of the Middle Miocene volcanics are covered by sediments and are known only from boreholes and geophysical studies.

In this study, we present new K/Ar ages on amphibole phenocrysts of volcaniclastic rocks from the Styrian Basin, Austria, as well as from the adjoining areas. Beside Lower Miocene shield volcanoes and Pliocene effusive alkaline volcanic rocks the eastern Styrian Basin hosts a number of phreatomagmatic tuff occurrences. The tuffs contain the well known mantle xenoliths and frequent amphibole phenocrysts. The new age data indicate that the Late Miocene phreatomagmatic volcanism started in Pontian and ended in Romanian. The complete interval of the youngest volcanism in the Styrian Basin covers 6 Ma, similar to the volcanism of the western part of the Pannonian Basin and the Nógrád/Novohrad area (West Carpathian).