



Ozone in the Mongolian Gobi desert: Large-scale concentrations and dry deposition

Franz X. Meixner (1), Thomas Behrendt (1,2,5), Michael Ermel (1), Nils Hempelmann (3), and Patrick Jöckel (4)

(1) Max Planck Institute for Chemistry, Biogeochemistry, Mainz, Germany (franz.meixner@mpic.de, +49-(0)6131-3056019),

(2) Institute of Geography, Gutenberg University Mainz, Mainz, Germany , (3) Climate Service Center (CSC), Hamburg,

Germany, (4) German Aerospace Center, DLR Oberpfaffenhofen, Institute for Atmospheric Physics, Wessling, Germany, (5)

Max Planck Graduate Center with Johannes Gutenberg-University Mainz GmbH, Mainz, Germany

For the first time, measurements of surface ozone mixing ratio have been performed from semi-arid steppe to arid/hyper-arid Mongolian Gobi desert. During 12-29 August 2009, ozone mixing ratio was continuously measured from a mobile platform (4x4 Furgon SUV). The survey (3060 km / 229171km²) started at the Mongolian capital Ulaan-Baatar (47.9582°N, 107.0190°E), heading to south-west (Echin Gol, 43.2586°N, 99.0255°E), eastward to Dalanzadgad (43.6061°N, 104.4445°E), and finally back to Ulaan-Baatar.

Ambient air was sampled (approx. 1 L/min) through a 4 m long PTFE-intake line along a forward facing boom mounted on the roof of a 4x4 Furgon SUV. Ozone mixing ratio has been measured by UV-spectroscopy using a mobile dual-cell ozone analyzer (model 205, 2BTechnologies, Boulder, U.S.A.). While ozone signals were measured every 5 seconds, 1 minute averages and standard deviations have been calculated on-line and stored into the data logger. The latter are used to identify and to discriminate against unrealistic low or high ozone mixing ratios which have been due to occasionally passing plumes of vehicle exhaust and/or biomass burning gases, as well as gasoline (at gas filling stations).

Even under desert conditions, the temporal behaviour of ozone mixing ratio was characterized by considerable and regular diel variations. Minimum mixing ratios (15-25 ppb) occurred early in the morning (approx. 06:00 local), when surface depletion of ozone (by dry deposition) can not be compensated by supply from the free troposphere due to thermodynamic stability of the nocturnal boundary layer. Late in the afternoon (approx. 17:00 local), under conditions of a turbulently well mixed convective boundary layer, maximum ozone mixing ratios (45-55 ppb) were reached. Daily amplitudes of the diel cycle of ozone mixing ratio were in the order of 30 ppb (steppe), 20 ppb (arid desert), to approx. 5 ppb (hyper-arid Gobi desert (Shargyn Gobi)).

Ozone surface measurements were compared to gridded ozone mixing ratios calculated by the global ECHAMMESSy model. Despite the model data's nature and the rather coarse spatial resolution (approx. 220 km) of the MESSy model, the agreement between (late afternoon) measured data and modelled data is surprisingly good.

Dry deposition velocity of ozone to steppe and desert soils will be estimated by application of the "Boundary Layer Mass Balance Technique" using the observed diel cycles of ozone mixing ratio and the temporal variation of the boundary layer height (a product of the MESSy model). Ozone dry deposition velocities estimated for the arid and hyper-arid (vegetation-free) regions of southern Mongolia will be compared to those deposition velocities which are known for pure ozone absorption to soils. Any differences are a strong indication for ozone sink mechanisms at the surface which are beyond physical absorption.