



## Timing and rates of long-term landscape evolution in Southern Argentina

Sebastian Kollenz (1), Ulrich A. Glasmacher (2), Eduardo A. Rossello (3), and Danny F. Stoeckli (4)

(1) Institute of Earth Sciences, University of Heidelberg, (2) Institute of Earth Sciences, University of Heidelberg, (3) Depto. de Ciencias Geológicas, FCEN, Universidad de Buenos Aires, (4) Department of Geological Sciences, University of Texas, Austin

The eastern Argentina South Atlantic passive continental margin is distinguished by a very flat topography. Out of the so called Pampean flat two mountain ranges are arising. These mountain ranges, the Sierras Australes and the Sierras Septentrionales, are located in the State of Buenos Aires south of the capital Buenos Aires. North of the Sierras Septentrionales the Salado basin is located. The Sierras Septentrionales and the Sierras Australes are also divided by a smaller intracratonic basin. Further in the South the Colorado basin is located. The Sierras Australes is a variscian fold belt originated by strong phases of metamorphism, but till now it is unclear by how many tectonic phases the area was influenced (Tomezzoli & Vilas, 1999). It consists of Proterozoic to Paleozoic rocks. The Sierras Septentrionales consists mainly of Precambrian crystalline rocks. The Precambrian sequences are overlain by younger Sediments (Cingolani, 2010). The aim is to understand the long-term landscape evolution of the area by quantifying erosion- and exhumation-rates and by dating ancient rock-uplift-events. Another goal is to find out how the opening of the south atlantic took effect on this region. To fulfill this goal, thermochronological techniques, such as fission-track dating and (U-Th-Sm)/He dating has been applied to samples from the region. Because there was no low- temperature thermochronology done in this area, both techniques were applied on apatites and zircons. Furthermore, numerical modeling of the cooling history has provided the data base for the quantification of the exhumation rates. The data-set show clusters of different ages which can be linked to tectonic activities during late Paleozoic times. Also the thermokinematic modeling is leading to new insights of the evolution of both mountain ranges and shows patterns of ongoing tectonic processes in this region. Calculated exhumation rates show also varying cooling histories and the influence of tectonics throughout the research area.

### References:

Tomezzoli, R. N. and Vilas, J. F. (1999): Palaeomagnetic constraints on the age of deformation of the Sierras Australes thrust and fold belt, Argentina. *Geophys. J. Int.* (1999) 138, 857–870.

Cingolani, C. A. (2010): The Tandilia System of Argentina as a southern extension of the Rio de la Plata craton: an overview, *Int. J. Earth. Sci. (Geol. Rundsch.)* (2011) 100, 221–242