Geophysical Research Abstracts Vol. 16, EGU2014-12550, 2014 EGU General Assembly 2014 © Author(s) 2014. CC Attribution 3.0 License.



The new diatom training set from the Polish Baltic coast and diatom-based transfer functions as a tool for understanding past changes in the southern Baltic coastal lakes

Monika Lutyńska (1), Grażyna Szpikowska (1), Michał Woszczyk (1), Anita Suchińska (2), Lubomira Burchardt (2), and Beata Messyasz (2)

(1) Adam Mickiewicz University, Institute of Geoecology and Geoinformation, Dziegielowa 27, 61-680 Poznan, Poland (lutynska@amu.edu.pl), (2) Adam Mickiewicz University, Department of Hydrobiology, Institute of Environmental Biology, Poland, Umultowska 89, 61-614 Poznan

The transfer function method has been developed as a useful tool for reconstruction of the past environmental changes. It is based on the assumption that the modern species, which ecological requirements are known, can be used to quantitative reconstructions of the past changes.

The aim of the study was to gather test sets and to build diatom-based transfer function, which can be used to reconstruct changes in the trophic state and salinity in the coastal lakes on the Polish Baltic coast. In the previous years there were several attempts made to reconstruct these parameters in lagoonal waters on the Baltic coasts in Germany, Denmark, Finland, Netherland, Sweden and Norway. But so far there is no diatom test set and transfer function was built for the Polish coastal lakes.

We sample diatoms form 12 lakes located along the polish Baltic coast. At the same time we monitor the physical-chemical conditions in the lakes, which includes lake water chemical composition (chlorides, phosphorous and sulphur), pH, salinity, conductivity, temperature, dissolved oxygen. We collect samples form the lakes as well as from the Baltic Sea and we analyse the whole phytoplankton composition, however the special focus in put on diatoms.

The results of the analysis show seasonal changes in the chemical and physical water properties. The diatom assemblage composition and species frequency also changed significantly.

This study is a contribution to the projects: NN 306 064 640 financed by National Science Centre, Poland and Virtual Institute ICLEA (Integrated Climate and Landscape Evolution Analysis) funded by the Helmholtz Association.