



Delta progradation in the wake of a large glacier surge: the Kuannersuit Sulluat delta, Disko, West Greenland

Verner Brandbyge Ernstsen (1), Marius Becker (2), and Thor Nygaard Markussen (1)

(1) Department of Geosciences and Natural Resource Management, University of Copenhagen, Denmark, (2) MARUM, University of Bremen, Germany

Beginning in 1995, a large outlet glacier of the Sermersauq Ice Cap on Disko Island surged 10.5 km downvalley to within 10 km of the head of the fjord, Kuannersuit Sulluat, reaching its maximum extent in summer 1999 before beginning to retreat (Gilbert et al., 2002). Bathymetric mapping of the delta front was carried out in summer 1999, at the maximum extent of the surged glacier, and again in summer 2013, i.e. 14 years after the initiated retreat of the outlet glacier. The location of the delta front in time is reconstructed from satellite images. Combining these analyses provides the potential for reconstructing volumetric delta progradation in response to the surging glacier event in the 1990s and the following retreat. The volumetric delta progradation will be compared with sediment transport measurements carried out upstream in the braided river during the period 1999-2001.

Acknowledgements

This work is funded by the Danish Council for Independent Research | Natural Sciences (Steno Grant 10-081102) and the German Research Foundation DFG-Research Center / Excellence Cluster “The Ocean in the Earth System”.

References

Gilbert R, Nielsen N, Möller H, Desloges JR, Rasch M (2002). Glacimarine sedimentation in Kangerdluk (Disko Fjord), West Greenland, in response to a surging glacier. *Marine Geology* 191: 1-18.