



## **Methane emission from Western Siberia derived from the integral methane balance in the troposphere for 2001 -2011**

Vasiliy Bogomolov (1), Victor Stepanenko (2), Igor Okladnikov (1), and Alexander Titov (1)

(1) Institute of Monitoring of Climatic and Ecological Systems SB RAS, Siberian center for Environmental Research and Training, Tomsk, Russian Federation (titov@scert.ru), (2) Moscow State University, Research Computing Center, Faculty of Geography

Although a number of estimates of methane emissions from large Earth regions have been proposed, based on either empirical or inverse transport modeling approaches, these emissions still remain highly uncertain motivating the development of new methods of surface flux assessment. In this study a new mathematical formulation for calculation of surface flux of methane, or any other gaseous component of the atmosphere, is developed. In it, the surface flux is retrieved from the integral balance of methane in bounded atmospheric domain. This balance includes the surface flux, the net advective flux through lateral boundaries of the atmospheric domain, methane sink due to oxidation by hydroxyl radical, and the rate of change of total methane amount in the domain.

Western Siberia, being one of the most prominent surface methane sources in Northern hemisphere, is used as a test region for the method. The components of methane balance are calculated using methane concentrations and wind speeds from MACC reanalysis at 1.125 deg. grid ([http://www.ecmwf.int/research/EU\\_projects/MACC](http://www.ecmwf.int/research/EU_projects/MACC)), and hydroxyl concentrations provided by (Spivakovskiy, 2000). The total methane flux from Western Siberia is thus assessed at 6 h resolution for 2001-2011. The time averaged total flux corresponds well to empirical estimates at diurnal and annual timescales. It may be anticipated that the method will be useful in close future when the satellite-retrieved methane concentration profiles will achieve high accuracy.