



## Recent advances in the Mercator-Ocean reanalysis system: Application to an Arctic configuration

Charles-Emmanuel Testut (1), Gilles Garric (1), Gregory Smith (2), and Youyu Lu (3)

(1) Mercator Ocean, R&D Assimilation, Ramonville Saint Agne, France (ctestut@mercator-ocean.fr), (2) Meteorological Research Division, EC, Dorval, Canada, (3) Bedford institute of Oceanography, Fisheries and Oceans Canada, Dartmouth, Canada

In the framework of the European project MyOcean, Mercator Ocean, the French operational oceanography center, is in charge of the development and of the production of real time analysis and forecasts and reanalysis for the global ocean. In this context, a global operational ocean system (model and data assimilation) provides, for instance, analyses and forecasts on a daily basis at the resolution of 1/12°. The operational systems are all based on the ocean and sea ice model NEMO and the multivariate data assimilation system SAM2 (Système d'Assimilation Mercator V2). The assimilation method is a reduced order Kalman filter based on SEEK formulation with bias correction scheme for temperature and salinity and an Incremental Analysis Update.

The strong need of a realistic description of the mean state and variability of the rapid changing Arctic Ocean and its adjacent seas over the last decades motivated the use of the Canadian Arctic Ocean and Nordic seas configuration (CREG). This dedicated configuration developed by the Canadian research teams has been coupled to the multivariate data assimilation system SAM2. The objectives of this Arctic platform is both to improve the sea ice assimilation method used in the Mercator Ocean and Canadian analysis and forecasting systems and to produce reanalysis over recent periods at lower numerical cost in order to prepare global higher resolution reanalysis. After a description of this Arctic reanalysis system, this poster presents firstly preliminary results on the abilities of this configuration to reproduce sea ice extent and volume interannual variability without assimilation over the last decades and, secondly, the impact of assimilating sea ice data on the sea ice cover with short hindcasts experiments.