



## **Vegetation pattern formation in semiarid systems induced by long-range competition in the absence of facilitation mechanisms.**

Ricardo Martinez-Garcia (1), Justin M. Calabrese (2), Emilio Hernandez-Garcia (1), and Cristobal Lopez (1)

(1) Institute for Cross-Disciplinary Physics and Complex Systems, UIB-CSIC, Palma de Mallorca, Spain (ricardo@ifisc.uib-csic.es), (2) Ecology Center, Smithsonian Conservation Biology Institute, Front Royal VA, USA.

Regular patterns and spatial organization of vegetation have been observed in many arid and semiarid ecosystems worldwide, covering a diverse range of plant taxa and soil types. A key common ingredient in these systems is that plant growth is severely limited by water availability, and thus plants likely compete strongly for water. The study of such patterns is especially interesting because their features may reveal much about the underlying physical and biological processes that generated them in addition to giving information on the characteristics of the ecosystem. It is possible, for instance, to infer their resilience against anthropogenic disturbances or climatic changes that could cause abrupt shifts in the system and lead it to a desert state.

Therefore much research has focused on identifying the underlying mechanisms that can produce spatial patterning in water-limited systems (Klausmeier, 1999). They are believed to arise from the interplay between long-range competition and facilitation processes acting at smaller distances (Borgogno et al., 2009). This combination of mechanisms is justified by arguing that water percolates more readily through the soil in vegetated areas (short range), and that plants compete for water resources over greater distances via long lateral roots (long range). However, recent studies have shown that even in the limit of local facilitation patterns may still appear (Martinez-Garcia et al., 2013).

In this work (Martinez-Garcia et al., 2013b), we show that, under rather general conditions, long-range competition alone is the minimal ingredient to shape gapped and stripped vegetation patterns typical of models that also account for facilitation in addition to competition. To this end we propose a simple, general model for the dynamics of vegetation, which includes only long-range competition between plants. Competition is introduced through a nonlocal term, where the kernel function quantifies the intensity of the interaction and its range. When the finite range of the competitive interaction is considered used kernel functions with a finite range, whose Fourier transform may have negative values, patterns emerge in the system. This is a rather general condition if we consider the finite length of the roots responsible of long-range competition for water in plant ecosystems. Therefore, our findings support the notion that, under fairly broad conditions, only competition is required for patterns to occur and suggest that the role of short-range facilitation mechanisms may not be as fundamental to pattern formation as has previously been thought.

### REFERENCES:

- C.A. Klausmeier, *Science*, 284, 1826-1828 (1999).
- F. Borgogno, P. D'Odorico, F. Laio and L. Ridolfi, *Reviews of Geophysics*, 4, RG1005 (2009).
- R. Martinez-Garcia, J.M. Calabrese, and C. Lopez, *Journal of Theoretical Biology*, 333, 156-165 (2013).
- R. Martinez-Garcia, J. M. Calabrese, E. Hernandez-Garcia, and C. Lopez, *Geophysical Research Letters*, 40, 6143-6147,(2013).