



Surface wave tomography of central and northern Europe from automated inter-station dispersion measurements

Riaz Ahmed Soomro (1), Christian Weidle (1), Sergei Lebedev (2), Luigia Cristiano (1), Jan Behrmann (3), and Thomas Meier (1)

(1) Institute of Geosciences, Kiel, SH, Germany(soomro@geophysik.uni-kiel.de), (2) School of Cosmic Physics, Geophysics Section, Dublin Institute for Advanced Studies, Dublin, Ireland, (3) Geomar Helmholtz Centre for Ocean Research, Kiel, Germany.

With the rapid growth in the number of seismic stations globally, automated analysis routines are inevitable to extract seismic observables from these large data volumes. For surface wave tomography, phase velocity dispersion curves of fundamental mode surface waves yield information on the isotropic as well as anisotropic structure of the crust and upper mantle. We have now developed an algorithm that enables automated, accurate measurement of inter-station phase velocity dispersion curves through cross correlation of vertical (Rayleigh wave) and transverse component (Love wave) seismograms.

After testing various parameters for automation of the procedure we find that the automated selection of segments of a given phase-velocity curve requires three parameters only, namely (1) difference of the measured phase velocity curve from a background model, (2) a smoothness constraint and (3) a length criterion. Furthermore, before averaging phase velocities for one path outliers are rejected. We performed rigorous tests to optimize the selection parameters. Interestingly, rough perturbations in the phase velocity curves do not bias the average phase velocities towards larger values. This indicates a rather random perturbation of the phase velocities by noise and complicated non-plane wave propagation.

We successfully applied the method to more than one hundred thousand inter station paths in central and northern Europe, involving more than one million cross correlations on 20 years of data from permanent networks. After inversion of the fundamental mode phase velocity dispersion curves for both Rayleigh and Love waves, we obtain high resolution anisotropic phase velocity maps for periods between 10 and 250 seconds with a lateral resolution between about 50 to 200 kilometers in Central Europe.

At shorter periods the North-German and Polish sedimentary basins are clearly imaged. At lower crustal depth the Alpine crustal root is characterized by low velocities. At longer periods a shallow asthenosphere beneath the regions of Cenozoic volcanism is found in Central Europe. The transition zone between the shallow Central European asthenosphere and the thick mantle lithosphere beneath the East European Craton along the Trans-European Suture Zone shows a distinct internal 3D structure. At depths between about 70 km to 100 km the strongest lateral heterogeneity is found in the region of the Elbe lineament, whereas at larger depths below about 120 km a very sharp boundary is present beneath the Tornquist-Teisseyre-Zone.

Azimuthal anisotropy at lower crustal depths across central Europe allows to differentiate between crustal and lithospheric domains created during the Variscan, Caledonian and Cadomian orogenies. Fast directions are interpreted as reflecting syn-orogenic deformation processes, as the fast directions of azimuthal anisotropy in the underlying asthenosphere are different.