Geophysical Research Abstracts Vol. 17, EGU2015-14417, 2015 EGU General Assembly 2015 © Author(s) 2015. CC Attribution 3.0 License.



Activity of faults observed in caves of the Eastern Alps

Ivo Baroň (1), Lukas Plan (1), Bernhard Grasemann (2), Ivanka Mitrovič (1,2), and Josef Stemberk (3) (1) Karst and Cave Research Group, Department of Geology and Paleontology, Natural History Museum, Burgring 7, 1030 Vienna, Austria (ivo.baron@nhm-wien.ac.at), (2) Department of Geodynamics and Sedimentology, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria, (3) Institute of Rock Structure and Mechanics ASCR, V Holešovičkách 94/41, 182 09 Prague, Czech Republic

Major recent tectonic process in the Eastern Alps involves the Neogene and Quaternary lateral extrusion of parts of the Eastern Alps towards the Pannonian Basin coeval with north-south shortening of the collision realm between the Adriatic Plate and the Bohemian Massif (European Plate). Within the framework of the FWF project "Speleotect" (2013-2017), we observe recent activity of the major fault systems of the Eastern Alps, such as the (1) Salzach-Ennstal-Mariazell-Puchberg (SEMP), (2) Mur-Mürz, (3) Periadriatic, (4) Lavanttal, and (5) Vienna Basin marginal Faults. Totally seven high-accuracy 3D crack-gauges TM71 with automated reading devices were installed in five selected karst caves with faults younger than the particular caves and correlated to one of these fault zones. The recorded micro-displacement events have been compared to known regional fault kinematics and to regional seismic activity (seismic data provided by the ZAMG). Already within the first year of observation, several micro displacement events were registered; these events sometimes revealed the same mechanisms as the geologically documented kinematics of the particular active faults, but in some cases performed completely opposite kinematics. These micro displacement events occurred in seismically rather quiet periods, however, usually about 1 – 10 days prior to local seismic events of different magnitudes (varying between ML 0.1 and 3.3). Further, in some cases gravitational mass movements were recorded that accompanied the tectonic moments.