Geophysical Research Abstracts Vol. 17, EGU2015-15902, 2015 EGU General Assembly 2015 © Author(s) 2015. CC Attribution 3.0 License.



Get immersed and black in fire ash world. Field Wildgeographers experience.

Paulo Pereira (1), Artemi Cerdà (2), Ieva Misiune (3), and Antonio Jordan (4)

(1) Environmental Management Centre, Mykolas Romeris University, Ateities g. 20, LT-08303 Vilnius, Lithuania, paulo@mruni.eu, (2) Soil Erosion and Degradation Research Group, Department of Geography, University of Valencia, Valencia, Spain. artemio.cerda@uv.es / www.soilerosion.eu, (3) Environmental Management Centre, Mykolas Romeris University, Ateities g. 20, LT-08303 Vilnius, Lithuania, ieva.misiune@gmail.com, (4) MED_Soil Research Group. Dep. of Crystallography, Mineralogy and Agricultural Chemistry, University of Seville, Spain. ajordan@us.es

Ash is the footprint left by the fire. This residue is very important to landscape recover and is an important source of soil protection after the fire. Get immersed in fire ash world is the same thing of get dirty and do not be recognizable in a fire affected area. To measure ash in the field we have to be careful where to step, how to do the experimental design, collect samples and find with accuracy the places measured before. A good methodology is needed in design field experiments, collect ash samples and monitoring ash evolution (Cerdà and Doerr, 2008; Bodi et al., 2014; Pereira and Ubeda, 2010; Pereira et al. 2011; 2012, 2013, 2014, 2015).

The objective of this work is to share with the Avatar world the methodologies used when wildgeographers get immersed in fire ash world, including:

1) Identify the best study area

2) Experimental design

3) Sample collection

4) Get dirty and have fun in the field

5) Laboratory, Statistical and spatial analysis

Acknoledgements

The author are thankful for the support of the projects POSTFIRE (Soil quality, erosion control and plant cover recovery under different post-fire management scenarios, CGL2013-47862-C2-1-R) funded by the Spanish Ministry of Economy and Competitiveness; GL2008-02879/BTE, LEDDRA 243857, RECARE (Preventing and Remediating Degradation of Soils in Europe Through Land Care, FP7-ENV-2013-TWO STAGE), funded by the European Commission; and for the COST action ES1306 (Connecteur: Connecting European Connectivity Research), FUEGORED (Spanish Network of Forest Fire Effects on Soils, http://grupo.us.es/fuegored/) and to Comissionat per a Universitats i Recerca del DIUE de la Generalitat de Catalunya.

References

Bodi, M., Martin, D.A., Santin, C., Balfour, V., Doerr, S.H., Pereira, P., Cerda, A., Mataix-Solera, J. (2014) Wildland fire ash: production, composition and eco-hydro-geomorphic effects. Earth-Science Reviews, 130, 103–127.

Cerdà, A. y Doerr, S.H. 2008. The effect of ash and needle cover on surface runoff and erosion in the immediate post-fire period. Catena, 74, 256-263. doi:10.1016/S0341-8162(02)00027-9

Pereira, P., Cerdà, A., Úbeda, X., Mataix-Solera, J. Arcenegui, V., Zavala, L. 2015. Modelling the impacts of wildfire on ash thickness in a short-term period, Land Degradation and Development, (In Press), DOI: 10.1002/ldr.2195

Pereira, P., Cerdà, A., Úbeda, X., Mataix-Solera, J., Jordan, A. Burguet, M. (2013) Spatial models for monitoring the spatio-temporal evolution of ashes after fire – a case study of a burnt grassland in Lithuania, Solid Earth, 4, 153-165.

Pereira, P., Úbeda, X. (2010) Spatial variation of heavy metals released from ashes after a wildfire, Journal of Environmental Engineering and Landscape Management, 18(1), 13-22.

Pereira, P., Ubeda, X., Martin, D. (2012) Fire severity effects on ash chemical composition and water-extractable elements, Geoderma, 191, 105-114.

Pereira, P., Úbeda, X., Martin, D., Mataix-Solera, J., Guerrero, C. (2011) Effects of a low prescribed fire in ash water soluble elements in a Cork Oak (Quercus suber) forest located in Northeast of Iberian Peninsula, Environmental Research, 111(2), 237–247.

Pereira, P., Úbeda, X., Martin, D., Mataix-Solera, J., Cerdà, A., Burguet, M. (2014) Wildfire effects on extractable elements in ash from a Pinus pinaster forest in Portugal, Hydrological Processes, 28, 3681–3690