



## **A multidisciplinary analysis to constrain exhumation and recent erosion history of the Tethyan Himalaya, based on apatite (U-Th-Sm)/He and cosmogenic nuclides dates from Central Nepal (Takkhola graben and the Mustang granite)**

Ruben Rosenkranz, Mohammad Sahragard Sohi, and Cornelia Spiegel

Department of Geosciences, University of Bremen, Bremen, Germany (rubenk@uni-bremen.de)

The exhumation of the Himalayan arc has been studied intensively throughout the last decades. For the Tethyan Himalaya, however, the youngest exhumation history is still unclear, mostly because of the lack of a significant apatite content in most Tethyan sediments (Crouzet et al. 2007). For this study we are using apatite (U-Th-Sm)/He thermochronology and cosmogenic nuclides for investigating exhumation and denudation of the Tethyan Himalaya back through time. Apatite (U-Th-Sm)/He thermochronology is sensitive to temperatures of  $\sim 40$  to  $85^{\circ}\text{C}$  and thus to movements within the upper  $\sim 1.5$  to  $3$  km of the earth's crust.

During a recent field campaign, we sampled the Mustang granite, that intrudes the Tethyan marine sediments and the Takkhola-Graben. The graben can be seen as an unusual southern part of the normal faulting system affecting the whole Tibetan Plateau (Colchen, 1999). The timing of the activation of these faulting is still highly debated. The syntectonic filling of the Takkhola-Mustag graben consists of Mio-Pliocene fluvio-lacustrine deposits (Garzione et al. 2003). These were described as containing significant amounts of apatite, derived from the past erosion of the Mustang granitic body (Adhikari and Wagerich, 2011). Being only up to  $1\text{ km}$  thick, a post depositional thermal resetting of the apatite (U-Th-Sm)/He system is unlikely, so that the (U-Th-Sm)/He dates of the sediments are expected to have retained their information regarding source area exhumation. We took several sand samples from the Kali Gandaki River draining the present-day exposure of the Mustang granite. Furthermore, we sampled different stratigraphic levels of the Mio-Pliocene sedimentary rocks, i.e. from the Tetang and Takkahola formation deposited between  $11$  and  $7$  Ma. This sampling approach will not only provide information about the youngest denudation history of the Mustang granite /Tethyan Himalaya, but will also reveal insights into its past denudation and changes of denudation rates through time for the region north to the South Tibetan Detachment.

It is our expectation to link the results to the tectonic behavior of the South Tibetan Detachment or to the newly recognized Western Nepal Fault System. Corroborating our results with other ages and diverse methods will provide a robust constraint on the exhumation and erosion history of the Tethyan Himalaya, as well as insights on the U-Th/He dating technique.

### References

Adhikari, B.R. and Wagerich, M., 2011. Provenance evolution of collapse graben fill in the Himalaya-The Miocene to Quaternary Thakkhola-Mustang graben (Nepal). *Sedimentary Geology*, 233(1), pp.1–14.

Blythe, A.E., Burbank, D.W., Carter, A., Schmidt, K., Putkonen, J., 2007. Plio-Quaternary exhumation history of the central Nepalese Himalaya: 1. Apatite and zircon fission track and apatite [U-Th]/He analyses. *Tectonics*, 26(3).

Crouzet, C. et al., 2007. Temperature and age constraints on the metamorphism of the Tethyan Himalaya in Central Nepal: A multidisciplinary approach. *Journal of Asian Earth Sciences*, 30(1), pp.113–130.

Garzione, C.N., DeCelles, P.G., Hokinson, D.G., Ojha, T.P., Upreti, B.N., 2003. East-west extension and Miocene environmental change in the southern Tibetan plateau: Thakkhola graben, central Nepal. *Geological Society of America Bulletin*, 115(1), pp.3–20.

Colchen, M., 1999. The Takkhola-Mustang graben in Nepal and the latee Cenozoic extension in the Higher Himalayas. *Journal of Asian Earth Sciences*, 17, pp. 683-702.