



‘Storm-flood’ coastal systems in relation to delta front-shoreface facies models: applications to Miocene successions in the Baram Delta Province, NW Borneo

Daniel Collins, Howard Johnson, and Peter Allison

Department of Earth Science and Engineering, Imperial College London, United Kingdom (dsc108@imperial.ac.uk)

Offshore sediment delivery requires flows to overcome the net landward bottom stress induced by fairweather shoaling waves. Storm waves are the dominant offshore transporting flow interpreted in basic models of clastic coasts, commonly generalised as a ‘shoreface’: the two-dimensional coarsening and shoaling upwards succession dominated by wave-formed sedimentary structures. Yet this model is just one end-member in a continuum of shoreface and delta front models, which reflect fundamental differences in shoreline processes, energy level and textural character.

We interpret abundant upward coarsening successions in the mid-late Miocene Belait Formation, eastern Baram Delta Province of NW Borneo, to reflect mixed-process storm- and river-flood influenced delta front deposits. High rates of sediment input attributed to combined fluvial and storm influence, is reflected in the pervasive 0.5-2 m-thick, very-fine grained sandstone beds. The abundance of swaley, to locally hummocky, cross-stratification, with exquisitely preserved mm-scale laminae, and internal discontinuity surfaces indicates repeated erosion and deposition during predominantly aggradational sedimentation. In addition, there is significant anisotropy within the swaley cross stratification, indicating a subordinate unidirectional flow acting with the inferred dominant oscillatory storm-generated component. Common mud drapes at the top of beds/bedsets and the observed gradients in ichnofauna diversity and abundance attest to a return to ambient, low-energy conditions. In some cases, angular carbonaceous material, including cm-scale wood clasts, indicate minimal reworking of fluvial-derived sediment. As large tropical storms invariably cause both increased fluvial discharge and sediment delivery (‘storm floods’), as well as increased wave energy, we propose a combined storm-dominated and fluvial-influenced delta front environment.

This is further supported by the size and frequency of erosional gutter casts that commonly define the bases of these beds. These features are uniquely abundant throughout the study area and display both exceptionally large dimensions, with thicknesses up to 2 m and widths >10 m, and a wide range of geometries. Isolated gutter casts, often up to 1-2 m deep, occur with steep to overhanging walls. The abundance, dimensions and wide-variability of gutter casts observed is unlike those documented in traditional wave-dominated shoreline successions and also attests to high-energy, erosive storm-flood flows with large sediment supply. We suggest the basinward progradation, coalescence of several small delta fronts, which may show updrift versus downdrift asymmetry, are responsible for a ‘ragged blanket’ of river-bypassed sand. Subsequent marine reworking may increase lateral sand continuity resulting in more typical ‘shoreface-like’ successions.