Geophysical Research Abstracts Vol. 17, EGU2015-7823, 2015 EGU General Assembly 2015 © Author(s) 2015. CC Attribution 3.0 License.



Status of hot spring water development in Korea 2014

Cholwoo Lee, Chan-Hee Park, and Yong-Chan Cho

KIGAM, Geothermal resources, Daejeon, Korea, Republic Of (cleanrain1116@gmail.com)

As of 2014, a number of Oncheon (hot spring) sites in Korea is 447. 207 sites among the total sites are located in the south-east part of Korea, because geothermal gradient in the region is higher than the average geothermal gradient of 25[U+2103]/km in Korea. The higher gradient leads to reduced drilling cost. 50% of the Oncheon sites are alkaline and their pH is higher than 8.5. pH for other 43% is between 7.5 and 8.5. As for the ion-type of Oncheon, 55% is are primarily the type of Na-HCO₃. 20% of Oncheon is the type of Na-Cl secondarily. 9% of Oncheon is the type of Na-SO4. While the well drilled deepest is 2,003 m from the surface, the average depth is 747 m. The highest temperature of hot spring is 78[U+2103], while the average temperature is 30[U+2103]. The area of Oncheon development was big until year 2000. Since then, the development area has shrunk to less than 20,000m2 to reduce risks for initial investment and stabilize the Oncheon management.

Keywords : Hot spring, Geothermal gradient, Temperature