Geophysical Research Abstracts Vol. 17, EGU2015-8376, 2015 EGU General Assembly 2015 © Author(s) 2015. CC Attribution 3.0 License.



Groundwater and surface water interaction in a basin surrounded by steep mountains, central Japan

Koichi Ikeda (1), Maki Tsujimura (2), Toshiaki Kaeriyama (3), and Takanori Nakano (4)

(1) Graduate School of Life and Environmental Sciences, University of Tsukuba, Ibaraki, Japan (s1330306@u.tsukuba.ac.jp), (2) Faculty of Life and Environmental Sciences, University of Tsukuba, Ibaraki, Japan (mktsuji@geoenv.tsukuba.ac.jp), (3) Industrial Economics and Construction Department, Ono city office, Fukui, Japan (kaeriyama.t@city.fukui-ono.lg.jp), (4) Center for Research Promotion, Research Institute for Humanity and Nature, Kyoto, Japan (nakanot@chikyu.ac.jp)

Mountainous headwaters and lower stream alluvial plains are important as water recharge and discharge areas from the view point of groundwater flow system. Especially, groundwater and surface water interaction is one of the most important processes to understand the total groundwater flow system from the mountain to the alluvial plain.

We performed tracer approach and hydrometric investigations in a basin with an area 948 square km surrounded by steep mountains with an altitude from 250m to 2060m, collected 258 groundwater samples and 112 surface water samples along four streams flowing in the basin. Also, Stable isotopes ratios of oxygen-18 (18O) and deuterium (D) and strontium (Sr) were determined on all water samples.

The 18O and D show distinctive values for each sub-basin affected by different average recharge altitudes among four sub-basins. Also, Sr isotope ratio shows the same trend as 18O and D affected by different geological covers in the recharge areas among four sub-basins.

The 18O, D and Sr isotope values of groundwater along some rivers in the middle stream region of the basin show close values as the rivers, and suggesting that direct recharge from the river to the shallow groundwater is predominant in that region. Also, a decreasing trend of discharge rate of the stream along the flow supports this idea of the groundwater and surface water interaction in the basin.