



Nutrient load can lead to enhanced CH₄ fluxes through changes in vegetation, peat surface elevation and water table depth in ombrotrophic bog

Sari Juutinen (1,2), Jill Bubier (3), Tuula Larmola (1,4), Elyn Humphreys (5), Sini Arnkil (1), Cameron Roy (6), and Tim Moore (6)

(1) Department of Forest Sciences, University of Helsinki, Helsinki, Finland (sari.juutinen@helsinki.fi; sini.arnkil@helsinki.fi), (2) Department of Environmental Sciences, University of Helsinki, Finland (sari.juutinen@helsinki.fi), (3) Environmental Studies Department, Mount Holyoke College, South Hadley, USA (jbubier@mtholyoke.edu), (4) Natural Resources Institute Finland, Vantaa, Finland (tuula.larmola@luke.fi), (5) Department of Geography and Environmental Studies, Carleton University, Ottawa, Canada (elyn.humphreys@carleton.ca), (6) Department of Geography, McGill University, Montreal, Canada (cameron.roy@mail.mcgill.ca; tim.moore@mcgill.ca)

Atmospheric nitrogen (N) deposition has led to nutrient enrichment in wetlands, particularly in temperate areas, affecting plant community composition, carbon (C) cycling, and microbial dynamics. It is vital to understand the temporal scales and mechanisms of the changes, because peatlands are long-term sinks of C, but sources of methane (CH₄), an important greenhouse gas. Rainwater fed (ombrotrophic) bogs are considered to be vulnerable to nutrient loading due to their natural nutrient poor status. We fertilized Mer Bleue Bog, a Sphagnum moss and evergreen shrub-dominated ombrotrophic bog near Ottawa, Ontario, now for 11-16 years with N (NO₃ NH₄) at 0.6, 3.2, and 6.4 g N m⁻² y⁻¹ (~5, 10 and 20 times ambient N deposition during summer months) with and without phosphorus (P) and potassium (K). Treatments were applied to triplicate plots (3 x 3 m) from May – August 2000-2015 and control plots received distilled water. We measured CH₄ fluxes with static chambers weekly from May to September 2015 and peat samples were incubated in laboratory to measure CH₄ production and consumption potentials. Methane fluxes at the site were generally low, but after 16 years, mean CH₄ emissions have increased and more than doubled in high nitrogen addition treatments if P and K input was also increased (3.2 and 6.4 g N m⁻²yr⁻¹ with PK), owing to drastic changes in vegetation and soil moisture. Vegetation changes include a loss of Sphagnum moss and introduction of new species, typical to minerogenic mires, which together with increased decomposition have led to decreased surface elevation and to higher water table level relative to the surface. The trajectories indicate that the N only treatments may result in similar responses, but only over longer time scales. Elevated atmospheric deposition of nutrients to peatlands may increase loss of C not only due to changes in CO₂ exchange but also due to enhanced CH₄ emissions in peatlands through a complex suite of feedbacks and interactions among vegetation, microclimate, and microbial processes. It is uncertain, however, how the vegetation change continues due to collapsing surface and higher water table levels, and how that will affect future CH₄ emissions and C balance.