



High-resolution sulfur isotopes in ice cores identify large stratospheric volcanic eruptions

Andrea Burke (1,2), Michael Sigl (3,4), Jess Adkins (2), Guillaume Paris (2), and Joe McConnell (4)

(1) Department of Earth and Environmental Sciences, University of St Andrews, St Andrews, UK (ab276@st-andrews.ac.uk), (2) Division of Geological and Planetary Sciences, Caltech, Pasadena, CA, USA, (3) Laboratory of Radiochemistry and Environmental Chemistry, Paul Scherrer Institut Villigen, Switzerland, (4) Desert Research Institute, Nevada System of Higher Education, Reno, NV, USA

The record of the volcanic forcing of climate over the past 2500 years is reconstructed primarily from sulfate concentrations in ice cores. Of particular interest are stratospheric eruptions, as these afford sulfate aerosols the longest residence time and largest dispersion in the atmosphere, and thus the greatest impact on radiative forcing. Identification of stratospheric eruptions currently relies on the successful matching of the same volcanic sulphate peak in ice cores from both the Northern and Southern hemispheres (a “bipolar event”). These are interpreted to reflect the global distribution of sulfur aerosols by the stratospheric winds. Despite its recent success, this method relies on precise and accurate dating of ice cores, in order to distinguish between a true ‘bipolar event’ and two separate eruptions that occurred in close temporal succession. Sulfur isotopes can be used to distinguish between these two scenarios since stratospheric sulfur aerosols are exposed to UV radiation which imparts a mass independent fractionation (Baroni et al., 2007). Mass independent fractionation of sulfate in ice cores thus offers a novel method of fingerprinting stratospheric eruptions, and thus refining the historic record of explosive volcanism and its forcing of climate. Here we present new high-resolution (sub-annual) sulfur isotope data from the Tunu Ice core in Greenland over seven eruptions. Sulfur isotopes were measured by MC-ICP-MS, which substantially reduces sample size requirements and allows high temporal resolution from a single ice core. We demonstrate the efficacy of the method on recent, well-known eruptions (including Pinatubo and Katmai/Novarupta), and then apply it to unidentified sulfate peaks, allowing us to identify new stratospheric eruptions.

Baroni, M., Thieme, M. H., Delmas, R. J., & Savarino, J. (2007). Mass-independent sulfur isotopic compositions in stratospheric volcanic eruptions. *Science*, 315(5808), 84–87. <http://doi.org/10.1126/science.1131754>