

## Accuracy assessment of gridded precipitation datasets in the Upper Indus Basin

Zubair Bashir (1), Adil Bashir (2), Asif Khan (3,4,5)

(1) Independent Researcher, Peshawar, Pakistan (zubairbashir96@yahoo.com), (2) City University, Department of Computer Science, Peshawar, Pakistan, (3) University of Cambridge, Department of Engineering, Cambridge, UK, (4) KPK, University of Engineering and Technology, Department of Engineering, Peshawar, Pakistan, (5) International Institute for Applied System Analysis, Austria

Accurate precipitation data are vital for hydro-climatic modelling and water resources assessments. Based on mass balance assessment, this study investigates the accuracy of thirteen widely used precipitation gridded datasets for sub-basins in the Upper Indus Basin (UIB) in the Himalayas-Karakoram-Hindukush (HKH) region. These datasets are: 1) Global Precipitation Climatology Project (GPCP), 2) Climate Prediction Centre (CPC) Merged Analysis of Precipitation (CMAP), 3) National Centers for Environmental Prediction (NCEP) / National Center for Atmospheric Research (NCAR), 4) Global Precipitation Climatology Centre (GPCC), 5) Climatic Research Unit (CRU), 6) Asian Precipitation Highly Resolved Observational Data Integration Towards Evaluation of Water Resources (APHRODITE), 7) Tropical Rainfall Measuring Mission (TRMM), 8) European Reanalysis (ERA) interim, 9) PRINCETON, 10) European Reanalysis-40 (ERA-40), 11) Willmott and Matsuura, 12) WATCH Forcing Data based on ERA interim (WFDEI), and 13) the Japanese 55-year Reanalysis (JRA-55) data. Precipitation accuracy and consistency was assessed by physical mass balance involving sum of annual measured flow, estimated actual evapotranspiration (average of 4 datasets), estimated glacier mass balance melt contribution (average of 4 datasets), and ground water recharge (average of 3 datasets), during 1999-2010. The accuracy assessment suggests that all tested precipitation datasets significantly under-estimate precipitation in the Karakoram sub-basins, except JRA-55 data. For the Hindu Kush and Himalayan sub-basins most datasets under-estimate precipitation except all Reanalysis datasets. The analysis indicates that for this large region with complicated terrain features and stark spatial precipitation gradients the reanalysis datasets have better consistency with flow measurements than datasets derived from records of only sparsely distributed climate stations in the HKH. Given the importance of the HKH region as the water tower of South Asia, our findings also suggest a revisit and scrutiny of hydro-climatic studies in the study region as to biases caused by the under-estimated gridded precipitation datasets.