

Anthropogenic perturbation of the global carbon cycle as a result of agricultural carbon erosion and burial

Zhengang Wang (1), Gerard Govers (2), Jed Kaplan (3), Thomas Hoffmann (4), Sebastian Doetterl (5), Johan Six (6), and Kristof Van Oost (1)

(1) UC Louvain, Louvain-la-neuve, Belgium (zhengang.wang@uclouvain.be), (2) Department of Earth and Environmental Sciences, K.U. Leuven, 3001 Heverlee, Belgium, (3) Institute of Earth Surface Dynamics, University of Lausanne, Geopolis, 1015 Lausanne, Switzerland, (4) Department of Geography, University Bonn, Meckenheimer Allee 166, DE-53115 Bonn, Germany, (5) Augsburg University, Institute of Geography, Augsburg, Germany, (6) Department of Environmental Systems Science, Swiss Federal Institute of Technology, ETH-Zurich, 8092 Zurich, Switzerland

Changes in terrestrial carbon storage exert a strong control over atmospheric CO_2 concentrations but the underlying mechanisms are not fully constrained. Anthropogenic land cover change is considered to represent an important carbon loss mechanism, but current assessments do not consider the associated acceleration of carbon erosion and burial in sediments. We evaluated the role of anthropogenic soil erosion and the resulting carbon fluxes between land and atmosphere from the onset of agriculture to the present day. We show, here, that agricultural erosion induced a significant cumulative net uptake of 198 ± 57 Pg carbon on terrestrial ecosystems. This erosion-induced soil carbon sink is estimated to have offset $74\pm21\%$ of carbon emissions. Since 1850, erosion fluxes have increased 3-fold. As a result, the erosion and lateral transfer of organic carbon in relation to human activities is an important driver of the global carbon cycle at millennial timescales.