Geophysical Research Abstracts Vol. 18, EGU2016-2264, 2016 EGU General Assembly 2016 © Author(s) 2016. CC Attribution 3.0 License.



METEOPOLE-FLUX: an observatory of terrestrial water, energy, and CO_2 fluxes in Toulouse

Jean-Christophe Calvet, Jean-Louis Roujean, Sibo Zhang, William Maurel, Bruno Piguet, Joël Barrié, Gilles Bouhours, Jacques Couzinier, Olivier Garrouste, Sandrine Girres, David Suquia, and Diane Tzanos CNRM-GAME (Météo-France, CNRS), Toulouse, France (jean-christophe.calvet@meteo.fr)

The METEOPOLE-FLUX project (http://www.cnrm.meteo.fr/spip.php?article874&lang=en) aims at monitoring a large suburban set-aside field in the city of Toulouse (43.572898 N, 1.374384 E). Since June 2012, these data contribute to the international effort to monitor terrestrial ecosystems (grasslands in particular), to the validation of land surface models, and to the near real time quality monitoring of operational weather forecast models. Various variables are monitored at a subhourly rate: wind speed, air temperature, air humidity, atmospheric pressure, precipitation, turbulent fluxes (H, LE, CO₂), downwelling and upwelling solar and infrared radiation, downwelling and upwelling PAR, fraction of diffuse incoming PAR, presence of water intercepted by vegetation (rain, dew), soil moisture profile, soil temperature profile, surface albedo, transmissivity of PAR in vegetation canopy. Moreover, local observations are performed using remote sensing techniques: infrared radiometry, GNSS reflectometry, and multi-band surface reflectometry using an aerosol photometer from the AERONET network. Destructive measurements of LAI, green/brown above-ground biomass, and necromass are performed twice a year. This site is characterized by a large fraction of gravels and stones in the soil, ranging from 17% to 35% in the top soil layer (down to 0.6 m), and peaking at 81% at 0.7 m. The impact of gravels and stones on thermal and moisture fluxes in the soil has not been much addressed in the past and is not represented in most land surface models. Their impact on the available water content for plant transpiration and plant growth is not much documented so far. The long term monitoring of this site will therefore improve the knowledge on land processes. The data will be used together with urban meteorological data to characterize the urban heat island. Finally, this site will be used for the CAL/VAL of various satellite products in conjunction with the SMOSMANIA soil moisture network (http://www.cnrm.meteo.fr/spip.php?article251&lang=en). The site will be presented together a first comparison of the ISBA land surface model with the observations.