

Influence of soil sampling approaches in the evaluation of soil organic carbon stocks under different land uses in a Mediterranean area

Rosa Francaviglia (1), Luca Doro (2), Luigi Ledda (2), Luis Parras-Alcántara (3), and Beatriz Lozano-García (3)

(1) Centro di Ricerca per lo Studio delle Relazioni tra Pianta e Suolo, Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria, Rome, Italy (rosa.francaviglia@entecra.it), (2) Università di Sassari, Dipartimento di Agraria, Sezione di Agronomia, Coltivazioni erbacee e Genetica, Italy, (3) University of Córdoba, Department of Agricultural Chemistry and Soil Science, Spain

Different approaches of soil sampling can provide significantly different estimates of soil organic carbon stocks (SOCs) (Parras-Alcántara et al., 2015a). Many studies have focused on SOC distribution only in the biologically active layers of topsoil, the IPCC carbon accounting method estimates the change in SOC storage for the top 30 cm of a soil profile, and indeed limited data are available for SOCs below this depth. Moreover, SOC estimates are more uncertain in areas with heterogeneous land uses and pedoclimatic conditions such as Mediterranean environments, which are more prone to land degradation due to SOC degradation and depletion and erosive processes (Muñoz-Rojas et al., 2015). Anyhow, the open question is whether soil should be sampled following the pedogenetic horizons with soil entire soil approach (ESP), or along fixed depth increments using the soil control section method (SCS) (Parras-Alcántara et al., 2015b). In addition, SOCs are often not adjusted for the soil volume occupied by coarse fragments as recommended by the IPCC Good Practice Guidance for LULUCF (IPCC, 2003) accordingly to the equation:

$$\text{SOCS} = \text{SOC} (\text{g kg}^{-1}) \times \text{bulk density} (\text{Mg m}^{-3}) \times \text{depth (m)} \times (1 - \text{coarse fragment}) \times 10.$$

The work deals with the comparison of SOCs using the ESP and SCS approaches, applied to a study area of northeastern Sardinia (Italy) under typical agro-silvo-pastoral systems (Francaviglia et al., 2014). The area lies within a hilly basin where elevation is in the range 275–340 m a.s.l., and slope ranges from 2–6% to 16–30%. The local climate is warm temperate with dry and hot summers, mean annual rainfall is 623 mm (range 367–811 mm) and mean annual temperature is 15.0°C (13.8–16.4°C). The area has the same soil type (Haplic Endoleptic Cambisols, Dystric) according to IUSS Working Group WRB (2006), and the following land uses with different levels of cropping intensification were compared: Tilled vineyards (Tv), No-tilled grassed vineyards (Ntgv), Hay crop (Hc), Pasture (P), Cork oak forest (Cof), former vineyards revegetated by Scrublands (Sfv), Mediterranean Maquis (Mmfv), and Helichrysum meadows (Hmfv). Average total SOCs were 128.0 and 140.6 Mg ha⁻¹ with the ESP and SCS approaches respectively if the coarse fraction is not included in the equation, 79.0 and 90.4 Mg ha⁻¹ when the coarse fraction is included. This indicates the importance to consider the coarse fraction when estimating SOC stocks, and an overestimation of SOCs when SCS sampling approach is adopted equal to about 16%.

References

- Francaviglia, R., Benedetti, A., Doro, L., Madrau, S., Ledda, S., 2014. Influence of land use on soil quality and stratification ratios under agro-silvo-pastoral Mediterranean management systems. *Agriculture, Ecosystems and Environment* 183, 86–92.
- Intergovernmental Panel on Climate Change (IPCC), 2003. In: Penman, J., Gytarsky, M., Hiraishi, T., Krug, T., Kruger, D., Pipatti, R., Buendia, L., Miwa, K., Ngara, T., Tanabe, K., Wagner, F. (Eds.), *Good Practice Guidance for Land Use, Land Use Change and Forestry*. IPCC/OECD/IEA/IGES, Hayama, Japan.
- IUSS Working Group WRB, 2006. World reference base for soil resources 2006. World Soil Resources Reports No. 103. FAO, Rome.
- Muñoz-Rojas, M., Doro, L., Ledda, L., Francaviglia, R., 2015. Application of CarboSOIL model to predict the effects of climate change on soil organic carbon stocks in agro-silvo-pastoral Mediterranean management systems. *Agriculture, Ecosystems and Environment* 202, 8–16.
- Parras-Alcántara, L., Lozano-García, B., Brevik, E. C., Cerdá, A., 2015a. Soil organic carbon stocks quantification in Mediterranean natural areas, a trade-off between entire soil profiles and soil control sections. *EGU General*

Assembly. Geophysical Research Abstracts Vol. 17, 2015-9865.

Parras-Alcántara, L., Lozano-García, B., Brevik, E., C., Cerdá, A., 2015b. Soil organic carbon stocks assessment in Mediterranean natural areas: A comparison of entire soil profiles and soil control sections. Journal of Environmental Management, 155, 219-228.