

The Mirdita Eastern Ophiolite, Albania: petrological evidence for MORB Mantle in the Tropoje peridotite massif?

Theodoros Ntaflos (1), Kujtim Onuzi (2), Christoph Hauzenberger (3), A. Hugh N. Rice (4), Peter Proßegger (1), Nazir Mekshiqi (2), and Gjon Kaza (2)

(1) University of Vienna, Dept. of Lithospheric Research, Austria (theodoros.ntaflos@univie.ac.at), (2) Institute of GeoSciences, Energy, Water and Environment, Tirana, Albania, (3) NAWI Graz Geocenter - Petrology & Geochemistry, Austria, (4) University of Vienna, Dept. of Geodynamics & Sedimentology, Austria

The Mirdita ophiolites in northern Albania are divided into the Western Mirdita Ophiolite (WMO), with MORB geochemical affinities, and the Eastern Mirdita Ophiolite (EMO), for which suprasubduction geochemical affinities have been reported. Ultramafic massifs in the WMO are often plagioclase bearing lherzolites/harzburgites whereas those in the EMO are strongly depleted spinel harzburgites.

The Tropoje ultramafic-mafic complex, which forms the northernmost part of the EMO, consists of spinel harzburgites, wherlites, orthopyroxenites and gabbros. The spinel harzburgites, which experienced variable degrees of serpentinization, are coarse- to medium-grained. One of the most striking features of the Tropoje ultramafic rocks is an outcrop of exceptionally fresh coarse- to medium-grained spinel harzburgites. While occasionally the peridotites exhibit a weak foliation, the prevalent texture is protogranular. Olivine is mainly coarse-grained (up to 6 mm size), typically showing kink-bands that frequently contain submicron-sized spinel exsolution lamellae. Both orthopyroxene and clinopyroxene, with grain size up to 3 mm and 1.5 mm, respectively, carry very thin exsolution lamellae of the other pyroxene. Spinel, up to 1 mm in size, is interstitial between boundaries of the associated silicates.

The mineral compositions of the spinel harzburgites indicate that they are strongly depleted in basaltic components. The rock-forming minerals (olivine, orthopyroxene and clinopyroxene) are all highly magnesian and chemically homogenous. The magnesium numbers ($Mg\# = 100 \times Mg / [Mg + Fe]$) for olivine and orthopyroxene are fairly homogeneous and vary within the narrow range of 90.9-91.6 and 91.5-91.7, respectively. The clinopyroxene is exceptionally highly magnesian, with $Mg\#$ ranging between 93.6 and 95.4. In both orthopyroxene and clinopyroxene, the Al_2O_3 contents range from 0.95 to 1.75 and from 0.62 to 2.27 wt%, respectively. Spinel shows a considerable variation in Al_2O_3 and Cr_2O_3 . The range in $Cr\#$ ($Cr\# = 100 \times Cr / [Cr + Al]$) is 51.7-69.2, but no compositional variations have been observed between core and rim.

The silicate minerals are strongly depleted in trace elements. LA-ICP-MS analyses of clinopyroxenes show that the LREE are below detection limit whereas the ratio of (Tb/Yb)_N range from 0.10 to 0.15 and Lu is around 0.5xPM.

Equilibrium temperatures calculated at a pressure of 1.5 GPa are relatively low, ranging between 620 and 830°C. The lowest temperature corresponds to the most residual peridotite, in which the clinopyroxene has the highest $Mg\#$ (95.4) and the spinel the highest $Cr\#$ (69.2).

The Tropoje ultramafic rocks have experienced high degrees of partial melting. Applying the method of Hellebrand (2001) to samples containing spinel with $Cr\# < 60$ yields around 20% partial melting, whereas samples with $Cr\# > 60$ suggest up to 30% partial melting.

The overall strong depletion in trace elements and prominent depletion of the LREE in clinopyroxenes as well as the absence of hydrous phases do not support a suprasubduction origin for the studied spinel harzburgites from Tropoje. Instead, the data suggest that our samples, from the Tropoje area are the residues of a MORB mantle after up to 30% partial melt extraction.