Geophysical Research Abstracts Vol. 19, EGU2017-12892, 2017 EGU General Assembly 2017 © Author(s) 2017. CC Attribution 3.0 License.



The CzeCOS Network

Kateřina Havránková, Klára Taufarová, Ladislav Šigut, Ryan McGloin, Manuel Acosta, Jiří Dušek , Lenka Krupková, Lenka Macálková-Mžourková, Marian Pavelka, Eva Dařenová, Shilpi Yadav, Vinh Nguyen, Carlos Guerra, Dalibor Janous, and Michal V. Marek

Global Change Research Institute CAS, Brno, Czech Republic

The Global Change Research Institute of the Czech Academy of Sciences (CzechGlobe) have established a wellequipped network of ecosystem stations, with modern instrumentation for eco-physiological, plant physiological and micrometeorological studies, and estimation of GHG emissions. The network of stations (CzeCOS) covers the main terrestrial ecosystems of the Czech Republic (young and old coniferous forest, deciduous forest, mixed floodplain forest, grassland, wetland and cropland). The ecosystem stations are equipped with eddy covariance systems, soil and stem chamber systems for CO_2 efflux and instruments for making micrometeorological measurements. The network enables detailed research to be conducted on topics such as: the carbon balance of different ecosystems, energy balance closure, the impact of current climate conditions on production and ecosystem disturbances during extreme weather conditions (drought, floods, winter storms, etc.) at regional, national and international scales. As a part of global networks (Fluxnet, ANAEe, ICOS), CzeCOS participates in evaluating and predicting environmental change and helps in the proposal of mitigation measures. Another important issue studied at some of the CzeCOS sites is the use of the eddy covariance method in sloping terrain in order to improve eddy covariance data processing for sites in this kind of terrain. Here we show specific results from the sites and outline the importance of the regional/national network for improving our knowledge about the exchange of matter and energy fluxes at different ecosystems.

This study was supported by the Ministry of Education, Youth and Sports of CR within the National Sustainability Program I (NPU I), grant number LO1415 and LD 15040.

Computational resources were provided by the CESNET LM2015042 and the CERIT Scientific Cloud LM2015085, provided under the programme "Projects of Large Research, Development, and Innovations Infrastructures".