

Analyzing scales of precipitation in general circulation models and observations

Nicholas Klingaman (1,2), Gill Martin (3), and Aurel Moise (4)

(1) National Centre for Atmospheric Science-Climate, Reading, UK (nicholas.klingaman@ncas.ac.uk), (2) Department of Meteorology, University of Reading, Reading, UK (nicholas.klingaman@ncas.ac.uk), (3) Met Office, Exeter, UK (gill.martin@metoffice.gov.uk), (4) Bureau of Meteorology, Melbourne, Australia (aurel.moise@bom.gov.au)

General circulation models (GCMs) have been criticized for failing to represent observed scales of precipitation, particularly in the tropics where simulated rainfall is often said to be too light, frequent and persistent. Previous assessments have used temporally or spatially averaged precipitation, which offers little actionable information for model developers, since the physics-dynamics interactions that produce precipitation occur at the native gridscale and timestep.

We introduce a set of diagnostics (ASoP1) to compare the spatial and temporal scales of precipitation across GCMs and observations, which can be applied to data ranging from the gridscale and timestep to regional and sub-monthly averages (Klingaman et al., 2017). When applied to data from ten GCMs, ASoP1 diagnostics reveal that far from the “dreary” persistent light rain implied by daily mean data, on the native timestep and gridscale most GCMs produce a broad range of intensities ($1\text{--}100 \text{ mm day}^{-1}$). Several GCMs, including the Met Office Unified Model (MetUM), show quasi-random behaviour that may alter the spectrum of atmospheric waves. Averaging to a common spatial ($\approx 600 \text{ km}$) or temporal (3 hr) resolution reduces inter-model variability, demonstrating that averaging masks intrinsic model behavior.

To further explore intermittent MetUM timestep precipitation and connect it to longer- and larger-scale biases, we analyze MetUM simulations at a range of horizontal resolutions, including $\approx 16 \text{ km}$ simulations with and without a deep convective parameterization (Martin et al., 2017). With parameterized convection, intermittency is largely insensitive to resolution and timestep length, as are larger- and longer-scale variability. Switching off the parameterization results in very persistent but very sporadic rainfall. On the $\approx 100 \text{ km}$ scale, the spectra of oceanic 3-hr and daily mean rainfall in the parameterized configurations agree well with satellite-derived rainfall estimates. At ≈ 10 -day scales, the spectra indicate a lack of intra-seasonal variability. Over tropical land, MetUM often underestimates daily mean rainfall (related to a poor representation of the diurnal cycle) and lacks sub-seasonal variability.

References

Klingaman NP, Martin GM and Moise AF, 2017: ASoP (v1.0): A set of methods for analyzing scales of precipitation in general circulation models. *Geosci. Model Dev.*, **10**, 57–83, doi:10.5194/gmd-10-57-2017.

Martin GM, Klingaman NP and Moise AF, 2017: Connecting spatial and temporal scales of tropical precipitation in observations and the MetUM-GA6. *Geosci. Model Dev.*, **10**, 105–126, doi:10.5194/gmd-10-105-2017.