



Assimilation of neural network soil moisture in land surface models

Nemesio Rodriguez-Fernandez (1), Patricia de Rosnay (2), Clement Albergel (3), Filipe Aires (4), Catherine Prigent (4), Yann Kerr (1), Philippe Richaume (1), Joaquin Muñoz-Sabater (2), and Matthias Drusch (5)

(1) CNRS, CESBIO, Toulouse cedex 9, France (nemesio.rodriguez@cesbio.cnes.fr), (2) ECMWF, Reading, United Kingdom, (3) Meteo France, Toulouse, France, (4) Observatoire de Paris, Paris, France, (5) ESTEC, European Space Agency, Noordwijk, The Netherlands

In this study a set of land surface data assimilation (DA) experiments making use of satellite derived soil moisture (SM) are presented. These experiments have two objectives: (1) to test the information content of satellite remote sensing of soil moisture for numerical weather prediction (NWP) models, and (2) to test a simplified assimilation of these data through the use of a Neural Network (NN) retrieval. Advanced Scatterometer (ASCAT) and Soil Moisture and Ocean Salinity (SMOS) data were used. The SMOS soil moisture dataset was obtained specifically for this project training a NN using SMOS brightness temperatures as input and using as reference for the training European Centre for Medium-Range Weather Forecasts (ECMWF) H-TESSEL SM fields. In this way, the SMOS NN SM dataset has a similar climatology to that of the model and it does not present a global bias with respect to the model. The DA experiments are computed using a surface-only Land Data Assimilation System (so-LDAS) based on the H-TESSEL land surface model. This system is very computationally efficient and allows to perform long surface assimilation experiments (one whole year, 2012). SMOS NN SM DA experiments are compared to ASCAT SM DA experiments. In both cases, experiments with and without 2 m air temperature and relative humidity DA are discussed using different observation errors for the ASCAT and SMOS datasets. Seasonal, geographical and soil-depth-related differences between the results of those experiments are presented and discussed. The different SM analysed fields are evaluated against a large number of in situ measurements of SM. On average, the SM analysis gives in general similar results to the model open loop with no assimilation even if significant differences can be seen for specific sites with in situ measurements. The sensitivity to observation errors to the SM dataset slightly differs depending on the networks of in situ measurements, however it is relatively low for the tests conducted here. Finally, the effect of the soil moisture analysis on the NWP is evaluated comparing experiments for different configurations of the system, with and without (Open Loop) soil moisture data assimilation. Assimilation of ASCAT soil moisture improves the forecast in the tropics and adds information with respect to the near surface conventional observations. In contrast, SMOS degrades the forecast in the Tropics in July-September. In the Southern hemisphere ASCAT degrades the forecast in July-September both alone and using 2m air temperature and relative humidity. On the other hand, experiments using SMOS (even without screen level variables) improve the forecast for all the seasons, in particular, in July-December. In the northern hemisphere both with ASCAT and SMOS, the experiments using 2m air temperature and relative humidity improve the forecast in April-September. SMOS alone has a significant positive effect in July-September for experiments with low observation error. Maps of the forecast skill with respect to the open loop experiment show that SMOS improves the forecast in North America and to a lesser extent in northern Asia for up to 72 hours.