

## Estimate of radiocaesium derived FNPP1 accident in the North Pacific Ocean

Yayoi Inomata (1), Michio Aoyama (2), Takaki Tsubono (3), Daisuke Tsumune (4), and Masatoshi Yamada (5)

(1) Institute of Nature and Environmental Technology, Kanazawa Univ., Kanazawa, Japan (yinomata@se.kanazawa-u.ac.jp), (2) Institute of Environmental Radioactivity, Fukushima Univ., Fukushima, Japan, (3) Environmental Science Research Laboratory, Central Research Institute of Electric Power Industry, Chiba, Japan, (4) Environmental Science Research Laboratory, Central Research Institute of Electric Power Industry, Chiba, Japan, (5) Institute of Radiation Emergency Medicine, Hirosaki Univ., Aomori, Japan

$^{134}\text{Cs}$  and  $^{137}\text{Cs}$  (radiocaesium) were released to the North Pacific Ocean by direct discharge and atmospheric deposition released from the TEPCO Fukushima Dai-ichi Nuclear Power Plant (FNPP1) accident in 2011. After the FNPP1 accident, measurements of  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  were conducted by many researches. However, those results are only snapshots in order to interpret the distribution and transport of the released radiocaesium on a basin scale. It is recognized that estimation of the total amount of released  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  is necessary to assess the radioecological impacts of their release on the environment. It was reported that the inventory of  $^{134}\text{Cs}$  or  $^{137}\text{Cs}$  on the North Pacific Ocean after the FNPP1 accident was 15.2-18.3 PBq based on the observations (Aoyama et al., 2016a),  $15.3 \pm 1.6$  PBq by OI analysis (Inomata et al., 2016),  $16.1 \pm 1.64$  PBq by global ocean model (Tsubono et al., 2016). These suggest that more than 75 % of the atmospheric-released radiocaesium (15.2-20.4 PBq; Aoyama et al., 2016a) were deposited on the North Pacific Ocean.

The radiocaesium from the atmospheric fallout and direct discharge were expected to mixing as well as diluting near the coastal region and transported eastward across the North Pacific Ocean in the surface layer. Furthermore, radiocaesium were rapidly mixed and penetrated into the subsurface water in the North Pacific Ocean in winter. It was revealed that these radiocaesium existed in the Subtropical Mode Water (STMW, Aoyama et al., 2016b; Kaeriyama et al., 2016) and Central Mode Water (CMW, Aoyama et al., 2016b), suggesting that mode water formation and subduction are efficient pathway for the transport of FNPP1 derived radiocaesium into the ocean interior within 1-year timescale. Kaeriyama et al. (2016) estimated the total amount of FNPP1 derived radiocaesium in the STMW was  $4.2 \pm 1.1$  PBq in October–November 2012. However, there is no estimation of the amount of radiocaesium in the CMW. Therefore, it is impossible to discuss about the mass balance of radiocaesium injected into the North Pacific Ocean. In this study, we conducted the optimum interpolation (OI) analysis to estimate the inventory of radiocaesium in the ocean interior as well as surface sweater by using the measured activities. Furthermore, transport speed of radiocaesium in the surface layer in the North Pacific Ocean were also estimated. The data used in this study were derived from all of the available data reported by such as the Tokyo Electric Power Company (TEPCO), the Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT), and voluntary cargo ships. The data analysis period was until December 2015 after the FNPP1 accident.

It was found that the radiocaesium across the North Pacific Ocean were reached to  $180^\circ\text{E}$  around  $40^\circ\text{N}$  latitude at July, 2012 by OI analysis. The transport speed was estimated to  $8.5 \text{ cm s}^{-1}$ . These were reached to the coastal site of America continent and the activities were increased after the year of 2014. The transport speed across  $70^\circ\text{W}$  ( $40^\circ\text{N}$  latitude) was decreased to  $5.2 \text{ cm s}^{-1}$ .

We estimated the inventory of radiocaesium in the surface seawater (depth; 0-100m) during the periods from August to December, 2012, based on the OI analysis. Amount of  $^{134}\text{Cs}$  inventory was estimated to 4.7 PBq with decay-corrected to 1 October 2012 (7.9 PBq at the time on 11 March 2011). (In the case of  $^{137}\text{Cs}$ , the inventory was estimated to 12.5 PBq with decay-corrected to 1 October 2012 and 13 PBq at the time on 11 March 2011 which includes pre-Fukushima  $^{137}\text{Cs}$  derived from atmospheric weapons test conducted in late 1950s and early 1960s). These correspond to 43-53% of the injected  $^{134}\text{Cs}$  in the North Pacific Ocean. It was reported that the  $4.2 \pm 1.1$  PBq of  $^{134}\text{Cs}$  were distributed in the STMW (Kaeriyama et al., 2016), which corresponds to 22-28% of the injected  $^{134}\text{Cs}$  in the North Pacific Ocean. Taking into account these estimation, FNPP1 derived radiocaesium existed in the CMW in the North Pacific Ocean would be about 3-6 PBq.

(References)

Aoyama, M., Hamajima, Y., Hult, M., Uematsu, M., Oka, E., . 2016.  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  in the North Pacific Ocean derived from the March 2011 TEPCO Fukushima Dai-Ichi Nuclear Power Plant accident, Japan. Part one: surface pathway and vertical distributions. *J. Oceanogr.* 72:53–65.

Aoyama, M., Kajino, M., Tanaka, T. Y., Sekiyama, T.T., Tsumune, D., Tsubono, T., Hamajima, Y., Inomata, Y., Gamo, T., .2016.  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  in the North Pacific Ocean derived from the March 2011 TEPCO Fukushima Daiichi Nuclear Power Plant accident, Japan. Part two: estimation of  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  inventories in the North Pacific Ocean. *J. Oceanogr.* 72:53–65.

Inomata, Y., Aoyama, M., Tsubono, T., Tsumune, D., Hirose, K. 2016. Spatial and temporal distributions of  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  derived from the TEPCOFukushima Daiichi nuclear power plant accident in the North Pacific Ocean by using optimal interpolation analysis. *Environ. Sci. Process. Impacts* 18:126–36.

Tsubono, T., Misumi. K., Tsumune, D., Bryan, F.O., Hirose, K., Aoyama, M. 2016. Evaluation of radioactive cesium impact from atmospheric deposition and direct release fluxes into the North Pacific from the Fukushima Daiichi nuclear power plant. *Deep-Sea Res. I.* 115:10–21.

Kaeriyama, H., Shimizu, Y., Setou, T., Kumamoto, Y., Okazaki, M., Ambe, D., Ono, T. 2016. Intrusion of Fukushima-derived radiocaesium into subsurface water due to formation of mode waters in the North Pacific. *Sci. Report.*, 6:22010 | DOI: 10.1038/srep22010.