

## Towards the improvement of model internal consistency using water isotope composition measurements in a glacierized basin

Zhihua He (1), Katy Unger-Shayesteh (1), Sergiy Vorogushyn (1), Abor Gafurov (1), Stephan Weise (2), Olga Kalashnikova (3), and Bruno Merz (1)

(1) GFZ German Research Centre for Geosciences, Section of Hydrology, Potsdam (Germany), (2) UFZ Helmholtz Centre for Environmental Research UFZ, Department Catchment Hydrology, Halle (Germany), (3) CAIAG Central Asian Institute of Applied Geosciences, Department Climate, Water and Natural Resources, Bishkek (Kyrgyzstan)

This study developed an approach to model the water stable isotope composition in streamflow in a glaicerized basin. The spatial variability of water isotope composition in precipitation was simulated in a Rayleigh fractionation procedure using spatial temperature and precipitation. The fractionation and mixing processes of water isotope compositions in various water sources were modeled in a glacio-hydrological model, associated with the simulation of runoff generation processes. Observed water isotope composition in streamflow was additionally used for the calibration of model parameters. The model internal consistency was investigated by the model performance in the reproduction of water isotope compositions addressed are three-fold: 1) How does the proposed approach perform in the reproduction of water isotope compositions of snow and glacier melt runoff? (3)What a degree the model internal consistency can be improved aided by the water isotope composition measurements? This study is one of the first attempts to simulate the water stable isotope composition in streamflow in glacierized basins. Our findings could shed lights on the application of water isotope composition measurements for model constraining in glacierized basins.