

The stress field and its role in the evolution of asthenosphere and lithosphere

Leszek Czechowski and Marek Grad

University of Warsaw, Institute of Geophysics, Faculty of Physics, Warsaw, Poland (lczech@op.pl)

Traditionally, the boundary between lithosphere and asthenosphere (LAB) is defined by a difference in response to stress: the lithosphere remains elastic or brittle, while the asthenosphere deforms viscously and accommodates strain through plastic deformation.

The lithosphere is a thermal boundary layer for the convection in the mantle. The rheology of rocks depends on many factors. However, the basic differences of lithosphere and asthenosphere properties could be explained as a result of the temperature and pressure. The effective viscosity of mantle is proportional to $C \exp(A/q)$, where q is the ratio (melting temperature/temperature), C and A are positive constants. The mantle is not molten, so $q > 1$. If the temperature is close to the melting temperature then q is close to 1 and effective viscosity is low (e.g. 1018 Pa s). This situation is observed in asthenosphere.

The temperature of the upper part of lithosphere is low (q is high) but temperature is increasing fast. In the mantle below LAB, the temperature gradient is low (could be close to the adiabatic one). The melting temperature is increasing with depth faster than true temperature. Hence, q and the viscosity reach minimum value just below LAB and are increasing with depth in the mantle below. It is a typical situation.

The tectonic processes could change this picture. The lithospheric plate could be placed in the mantle below another plate. Distribution q in such a case could have two minima, so two asthenospheric layers could be formed.

Another important factor determining rheological properties is a stress tensor T . Generally viscosity is proportional to the power of the invariant of the stress tensor: $I(T)^{(1-n)}$. For $n=1$ the viscosity does not depend on stress (i.e. Newtonian rheology), for true mantle n is probably in the range from 3 to 5.

We investigate the processes of formation and evolution of low viscosity layers ("asthenospheric layers") in the upper mantle. The time scale of the temperature changes is of the order of 10 - 20 Myr. The characteristic time of stress changes could be much shorter depending on tectonic processes. In a relatively short time (below 1 Myr) the pattern the viscosity distribution and velocity gradient could change substantially.

Using results from deep seismic sounding and surface wave tomography we have found that below some regions there are structures in the mantle that could be a forming/vanishing low viscosity layers. Reflectors in the lower lithosphere are observed beneath Trans-European Suture Zone between Precambrian and Palaeozoic platforms. In a thick Baltic Shield lithosphere (200 km or more) low velocity zones and seismic reflectors are observed in the depth range 60-100 km, which could be interpreted as mechanical low V_p velocity zones, in contrast to thermal velocity zone in the asthenosphere.

Acknowledgments: This work was partially supported by the National Science Centre (grant 2011/01/B/ST10/06653).